SOIL FERTILITY STATUS OF GUDAMALANI TALUKA OF BARMER DISTRICT OF RAJASTHAN

Ganga Ram Mali, Turfan Khan, Pradeep Pagaria* L.R. Choudhary and H.D. Choudhary

Krishi Vigyan Kendra, Barmer II- 344031 (Rajasthan), Agriculture university, Jodhpur

Received-09.06.2020, Revised-26.06.2020

Abstract: Within a soil, nutrient variability exists depending upon the hydrological properties of the soil and cropping system. In the present study 53 soil samples were collected from 9 gram panchayats and were analysed. The soil samples were collected from Pearlmillet-Cumin cropping sequence. Analysis of soil samples revealed that 100 per cent soil samples were deficient in available nitrogen and organic matter content, while all samples were medium range in P and K, respectively. Among the micronutrients tested copper and manganese were in sufficient range while iron and zinc were deficient in soil.

Keywords: Soil fertility, Nitrogen, Phosphorus, Organic matter, Micronutrients, Analysis

REFERENCES

Jackson, M.L. (1973). Soil Chemical Analysis, Prentice Hall of India Pvt. Ltd., New Delhi, p. 38-56. Lindsay, W. L. and Norvell, W. A. (1978). Development of DTPA soil test for zinc, iron, manganese and coper. *Soil Science Society America Journal* **42**: 421-428.

Olsen, S.R., Cole, C.V., Watanabe, F.S. and Dean, L.A. (1954). Estimation of available phosphorus in

soil by extraction with sodium bicarbonate. *Circular* 939, USDA, Washington, DC, USA.

Subbiah, B.V. and Asija, G.L. (1956). A rapid procedure for the estimation of available nitrogen in soils. *Current Science* **25**(7): 259-260.

Walkley, A. J. and Black, I. A. (1934). Estimation of soil organic carbon by the chromic acid titration method. *Soil Science* **37**: 29-38.