

EXISTING PRODUCTION PATTERNS AMONG THE MAIZE GROWERS

P.K. Netam*, Basanti Netam and Virendra Kumar Painkra

¹Department of Agricultural Extension CARS, Kanker, IGKV, Raipur, Chhattisgarh²Department of Senior Agriculture Development Officer, Dhamtari, Chhattisgarh³Department of Agricultural Extension CoA, IGKV, Raipur, ChhattisgarhEmail: pknetam49@gmail.com

Received-01.04.2020, Revised-22.04.2020

Abstracts: This investigation was carried out in three district of Bastar plateau of Chhattisgarh State to assess the level of existing production pattern among the respondents. 270 farmers were considering as respondents for this study. Respondents were interviewed through personal interview. Collected data were analyzed with the help of suitable statistical methods. The analysis of the results showed that major crop prevailed in *Kharif* season among different respondents was rice followed by maize, while predominant crop in *Rabi* season was maize covering 64.44 percent area.

Keywords: Production pattern, Area, Productivity, Economic assessment

REFERENCES

Chhattisgarh (2014). Annual statistics report.

CIMMYT (2005). Maize in India: production systems, constraints, and research priorities.

FAOSTAT (2012). Statistical databases and data sets of the food and agriculture organization of the United Nations.

FARA (2009). Pattern of change in maize production in Africa: implication for policy development. Ministerial policy brief services, No.3 December 2009 Accra, Ghana: forum for Agricultural Research in Africa (FARA).

Gecho, Yishak and Punjabi, N.K. (2011). Determination of adoption of improved Maize technology in Damot Gale, Wolaita, Ethiopia. *Raj. J. Ext. Edu.*, 19: 1-9.

Gupta, Km. Saroj and Gyanpur, S.R.N. (2012). Sustainability of scientific maize cultivation practice in Uttar Pradesh, India. *Journal of Agricultural Technology*. 8 (3): 1089-1098.

Kumari, Sunita, Sharma, F.L. and Nidhi (2017). Study of profile characteristics of wheat and maize

growers in Udaipur District of Rajasthan. *IMPACT: International Journal of Research in Applied, Natural and Social Sciences*, 5(2): 1-12.

Langade, D. M., Shahi, J.P., Agrawal, V. K. and Sharma, A. (2013). Maize as emerging source of oil in india: an overview. *Maydica*, 58(3/4): 224-230.

Nyangena, Wilfred and Juma, Ogada Maurice (2014). Impact of Improved Farm Technologies on Yields: The case of improved Maize Varieties and Inorganic fertilizer in Kenya. *Environment for Development*.

R. Cox (1956). Control of helminthosporium turcicum blight disease of sweet corn in South Florida. *Phytopathology*, 5: 68-70.

Singha, K. and Chakravorty, A. (2013). Crop diversification in India: a study of maize cultivation in Karnataka. *Scientific Journal of Review*, 2(1): 1-10.

Wokabi, S. M. (1998). Sustainability of maize production in Kenya. Kenya Agricultural Research Institute, Nairobi, Kenya. p. 2.