## EFFECT OF WATER STRESS ON YIELD COMPONENTS OF IRANIAN WHEAT LANDRACES UNDER IRRIGATED, RESTRICTED IRRIGATED AND RAIN-FED CONDITIONS

## Amandeep Kaur\* and Rashpal Singh Sarlach<sup>2</sup>

<sup>1</sup>Department of Botany, Punjab Agricultural University, Ludhiana 141004 <sup>2</sup>Department of Plant Breeding & Genetics, Punjab Agricultural University, Ludhiana,141004 Email:deepaman3305@gmail.com

## Received-03.06.2019, Revised-25.06.2019

**Abstract:** Drought stress is one of the most important environmental factor retard the formation of the yield components which are most actively developing at the time of stress which results in 17% to70% of grain yield. The aim of present study was to investigate the effect of water stress on morphological traits of 27 Iranian landraces along with commercial relevant checks under irrigated, restricted irrigated and Rain-fed conditions. These lines were selected on the basis of minimum reduction of vigor index under water stress induced by Polyethylene glycol (6000) as compared to control lines. A field experiment was carried out at experimental area of Department of Plant Breeding & Genetics, Punjab Agricultural University Ludhiana, Punjab during 2016 with three replications. On the basis of performance of Iranian lines under field conditions these three lines IWA 8600064, IWA 8600179 and PETTERSON ML68-10 are considered as water stress tolerant.

Keywords: Drought stress, Iranian landraces, Yield parameters

## REFERENCES

Kaur, A., Sarlach, R.S., Sharma, A. and Bains, N.S. (2018). Identification of drought tolerant Iranian wheat landraces under water stress conditions .Vegetos 31: 68-73.

**Bilal, M., Rashid, R.M., Rehman, S.U., Iqbal, F., Ahmed, J., Abid, M.A., Ahmed, Z. and Hayat, A.** (2015). Evaluation of wheat genotypes for drought tolerance. J G

Kheiralla, K.H., Ismai, A.A. and. El-Negar, G.R. (1997). Drought tolerance and stability of some spring wheatcultivars. Assiut J. Agriculture Sciences 28: 75-80.

Kilic, H. and Yağbasanlar, T. (2010). The effect of drought stress on grain yield, yield components and some quality traits of durum wheat (*Triticum turgidum*) cultivars. Notulae Botanicae Horti Agrobotanici Cluj-Napoca 38: 164-70.

**Lobell, D. B. and Gourdji, S. M.** (2012). The influence of climate change on global crop productivity. Plant Physiology 160: 1686–1697

Porter, J. R. and Semenov, M. A. (2005). Crop responses to climatic variation. Philosophical

Transactions of the Royal Society B: Biological Sciences 360: 2021–2035.

Saini, H.S. and Westgate, M.E. (1999). Reproductive development in grain crops during drought. Advance Agron 68: 59-96.

**Shiferaw, B. et al.** (2013). Crops that feed the world . Past successes and future challenges to the role played by wheat in global food security. Food Security 5:291–317.

**Stone, P.J. and Nicholas, M. E.** (1995). Comparison of sudden heat stress with gradual exposure to high temperature during grain filling in two wheat varieties differing in heat tolerance. I Grain growth. Aust. Journal of Plant Physiol., 22: 935-944.

**Zhang, H. and Oweis, T.** (1999). Water yield relation and optimal irrigation scheduling of wheat in the Mediterranean region. Agric. Water Management., 38:195-211.

\*Corresponding Author

Journal of Plant Development Sciences Vol. 11(6): 315-322. 2019