PESTICIDES APPLIATIONS AND THEIR PERSISTANCE IN VEGETABLES

Ankush Sheoran* and Rohit Rana

Department of Chemistry, Panjab University, Chandigarh Email: <u>ankushsheoran16@gmail.com</u>

Received-27.10.2017, Revised-16.11.2017

Abstract: Although pesticide application is an essential component of modern crop production technology, however, in recent years, contamination of food commodities especially the vegetables, with trace amounts of these chemicals has become a growing issue for the general population. Unfortunately the debate has restricted to the quality standards and norms pertaining to drinking water and foods at the point of consumption. But more fundamental source of contamination of natural resources including food products with chemical pesticide residues at the farming level is of serious concern, because of their persistence or presence and often ignored. Without solving this problem at basic level, setting down the quality standards at the consumption level is not going to solve the problem, especially in developing countries like India where enforcement of rules and regulations is very weak or absent. Although some pesticides application have banned but still their residues are being reported in food products and natural resources like soil and water due to its high persistence and low biodegradability. Therefore, it is essential to create awareness amongst consumers and producers about health risks and production practices like non-pesticidal management activities should be encouraged for healthy and safe environment.

Keyword: Pesticides application, Vegetable crops, Drinking water, Health hazards

REFERENCES

Islam, S., Afrin, N., Hossain, M.S., Nahar, N., Mosihuzzaman, M. and Mamun, M.I.R. (2009). Analysis of some pesticide residues in cauliflower by high performance liquid chromatography. American Journal of Environmental Science 5(3):325-329.

Debayle, D., Dessalces, G., Florence, M. and Loustalot, G. (2008). Multiresidue analysis of traces of pesticides and antibiotics in honey by HPLC-MS-MS, Analytical and bio-analytical chemistry 391(3): 1011-1020.

Abhilash, P.C., Singh, V. and Singh, N. (2009). Simplified determination of combined residues of lindane and HCH isomers in vegetables , fruits , wheat, pulses and medicinal plants by matrix solidphase dispersion (MSPD) followed by GC-ECD. *Food Chemistry*. 113(1): 267-271.

Akhtar, S.A. (2009). Dissipation and transloction of propineb in tomato. *Pesticide Research Journal* 22 (2): 145-147.

APEDA (Agricultural and Processed Food Products Export Development Authority) (2015). Fresh Fruits & Vegetables.Agricultural and Processed Food Products Export Development Authority (APEDA), Ministry of Commerce andIndustry, Government of India.

http://www.apeda.gov.in/apedawebsite/six_head_pro duct/FFV.htm.

Arora, S. and Singh, D.K. (2004). Determined of Chlorpyriphos and Cypermethrin residues in/on okra. *Pesticide Research Journal* 16 (2): 68-70.

Baker, J.L. *et al.*, (1994). Aquatic dialogue group: Pesticide risk assessment and mitigation, SETAC Press, Florida, pp. 188.

Bhanti, M. and Taneja, A. (2005). Monitoring of Organochlorine Pesticide Residues in Summer and Winter Vegetables from Agra, India – A case study. *Corresponding Author

Environmental Monitoring and Assessment. 110(1-3): 341-346.

Bhattacherjee, A.K. and Pandey, B.K. (2010). Dissipation of carbendazim in mango in pre and post harvest treatments. *Journal of Plant Protection science* 2 (1): 65-70.

Caldas, E.D., Conceiçao, M.H., Miranda, M.C.C., de Souza, L.C.K. and Lima, J.F. (2001). Determination of dithiocarbamate fungicide residue in food by spectrophotometric method using a vertical disulfide reaction system. *Journal of Agricultural and Food Chemistry* 49(10): 4521-4525. Charan, P.D., Ali, S.F., Yati K. and Sharma, K.C. (2010). Monitoring of pesticide residues in farmgate vegetables of central Aravalli region of Western India. *American-Eurasian Journal of Agricultural & Environmental Sciences* 7(3): 255-258.

Clark, J.M. and H. Ohkawa (eds.). (2003). Environmental fate and safety management of agrochemicals, ACS Symposium Series 899. ACS, Washington D.C., pp. 357.

Dhaliwal, G.S. (2009). An Outline of Entomology, Kalyani Publishers, Ludhiana.

Dikshit, A. K. and Mishra, S. S. (1985). Residues of endosulfan and carbaryl in potato tubers at harvest and after storage. *Indian Journal of Plant Protection* 13: 105-108.

Freemark, K. and Boutin, C. (1994). Non-target plant risk assessment for pesticide registration. *Environmental Management* 18(6): 841-854.

Gajbhiye, V.T., Kumar, R., Gupta, R.K. and Kalpana (2000). Translocation and Persistence of Imidacloprid in Tomato. *Pesticide Research Journal* 12 (1): 127-129.

Gupta, P.K. (2004). Pesticide exposure—Indian scene. *Toxicology* 198: 83–90.

Gupta, S., Sharma, R.K., Gupta, R.K., Sinha, S.R., Singh, R. and Gajbhiye, V.T. (2008).

Journal of Plant Development Sciences Vol. 9 (11): 977-981. 2017

Persistence of new insecticides and their efficacy against insect pests of okra. *Bulletin Environmental Contamination and Toxicology* 8: 1-9.

Janghel, E.K., Rai, J.K., Rai, M.K. and Gupta, V.K. (2007). A new sensitive spectrophotometric determination of cypermethrin insecticide in environmental and biological samples. *Journal of the Brazilian Chemical Society* 18(3): 590-594.

Kalara R.L. (2003). Pesticide residues in vegetable. In Proceedings of symposium on risk assessment of pesticide residues in water and food (pp. A1-9).By ILSI Washington DC, ITRC Lucknow and ICMR, New Delhi, India on October 28-29.

Karanth, N.G.K. (2002). Challenges of limiting pesticide residues in fresh vegetables: the Indian experience. In: Hanak, E.E., P. Boutrif. and M.P. Fabre. (Eds.), Food Safety Management in Developing Countries. CIRAD-FAO, Montpellier, France, pp. 11-13.

Kumari, B. (2008). Effect of household processing on reduction of pesticide residues in vegetables. *ARPN Journal of Agricultural and Biological Science* 3 (4): 46-51.

Ministry of Agriculture. (2015). Agricultural Statistics at a Glance 2014. Ministry of Agriculture, Government of India. Oxford University Press, New Delhi, India

Mohapatra,S., Sharma, D. and Rekha, A. (2006). Residues of Lambda cyhalothrin in/on Acid Lime (*Citrus aurantifolis*). *Pesticide Residue Journal* 18 (1): 74-75.

Mukherjee, I. and Gopal, M. (2003). Pesticide residues in vegetable. In Proceedings of symposium on risk assessment of pesticide residues in water and food (pp. A1-8).By ILSI Washington DC, ITRC Lucknow and ICMR, New Delhi, India on October 28-29.

Newsome, W.H., Doucet, J., Davies, D. and Sun, W.F. (2000). Pesticide residue in the Canadian market basket survey-1992 to 1996. *Food Additives and Contaminants* 17(10): 847-854.

Nishant, N. and Upadhyay, R. (2016). Presence of pesticide residue in vegetable crops: A review. *Agricultural Reviews* 37(3): 2016: 173-185.

Parveen, Z. and Masud, S.Z. (2003). Monitoring of pesticide residue in human milk. *Pakistan Journal of Science and Industrial Research* 46: 43-46.

Patil, V.B. and Shingare, M.S. (1994). Thin-layer chromatographic spray reagent for the screening of biological materials for the presence of carbaryl. *Analyst*, 119(3): 415-416.

Peshin, R., Kranthi, K.R. and Sharma, R. (2014). Pesticide use and experiences with integrated pest management programsand Bt cotton in India. In: Peshin, R., Pimentel, D. (Eds.), Integrated Pest Management Experiences with Implementation: Global Overview, vol. 4. Springer, Dordrecht, The Netherlands, pp. 255-268.

RangaRao, G.V., Sahrawat, K.L., Srinivasa, R.C., Binitha, D., Reddy, K.K. and Bharath, B.S. (2009). Insecticide residues invegetable crops grown in Kothapalli Watershed, Andhra Pradesh, India: A case Study. *Indian Journal of Dryland Agricultural Research & Development* 24: 21-27.

Rathore, H.S. and Begum, T. (1993). Thin-layer chromatographic behaviour of carbamate pesticides and related compound. *Journal of Chromatography A*, 643(1): 321-329.

Reddy, J.D., Rao, N.B. and Sultan, A.M. (2000). Insecticide Residues in market samples of grape berries. *Pestology* 16 (9):17-22.

Sardana, H.R. (2001). Integrated Pest Management in Vegetables, In: Training Manual on IPM for Zonal Agricultural Research Stations, pp. 105-118.

Sharma, I.D., Dubey, J.K. and Patyal, S.K. (2006). Persistence of Fenazaquin in Apple Fruits and Soil. *Pesticide Research Journal* 18 (1): 79-81.

Tan, G.H. and Soo M. T. (2001). Multiresidue analysis of pesticides in vegetables using liquid chromatography with atmospheric pressure ionization mass spectrometry (LC-API-MS) and a heated nebulizer. *Malaysian J. of Chemistry.* 3(1): 1-12.

Thanh, D.N., Ji, E.Y., Dae, M.L. and Gae, H.L. (2008). A multiresidue method for the determination of 107 pesticides in cabbage and radish using QuEChERS sample preparation method and gas chromatography mass spectrometry. *Food Chemistry*, 110(1): 207-213.

Vijayalakshmi, K. (2002). Dissipation of quinalphos on mango (*Mangifera indica*) fruit. *Tropical Agriculture (Trinidad)* 78 (1): 39-42.