

MOLECULAR METHODS FOR PLANT TAXONOMY

Vinay M. Raole* and Nagesh Chirumamilla

Department of Botany, Faculty of Science
 The Maharaja Sayajirao University of Baroda, Vadodara-390002
 Email: vinaysar@rediffmail.com

Received-02.08.2017, Revised-26.08.2017

Abstract: Molecular systematics is the use of molecular genetics to study the evolution of relationships among individuals and species. The goal of systematic studies is to provide insight into the history of groups of organisms and the evolutionary processes that create diversity among species. There are two separate tasks to which DNA specificity is currently being applied. First one DNA data used to distinguish between species which is equivalent to species identification and the second one to discover new species. The aim of this review is to present the techniques that are available to a taxonomist to complement the conventional field methods of identification and delineation of plant species.

Keywords: Genetic diversity, PCR, AFLP, DNA Barcode

REFERENCES

- Ali, I.F., D.B. Nelae and K.A. Marshall** (1991). Chloroplast DNA restriction fragment length polymorphisms in *Sequoia sempervirens* D. Don Endl., *Pseudotsuga menziesii* (Mirb). Franco, *Claocedrus decurrens* (Tott.) and *Pinus taeda* L. Theor. Appl. Genet. 81: 83-89.
- Alvarez , I.A. and Wendel, J.F.** (2003). Ribosomal ITS sequences and plant phylogenetic inference. Mol Phylogen Evol 29:417–434
- Alves, R.J.V. and Machado, M.D.** (2007). Is classical taxonomy obsolete? Taxon 56:287–288
- Bark, O.H. and M.J. Hayey** (1995). Similarities and relationships among populations of the bulb onion as estimated by RFLPs. Theor. Appl. Genet. 90:407-414.
- Besse, P., M. Seguin, P. Lebrun, M.H. Chevallier, D. Nicolas and C. Lanauad** (1994). Genetic diversity among wild and cultivated populations of *Hevea brasiliensis* assessed by nuclear RFLP analysis. Theor. Appl. Genet. 88:199-207.
- Bhargava, A., Fuentes, F.F.** (2010). Mutational dynamics of microsatellites. Mol Biotechnol 44:250–266
- Braslavsky, I.; Hebert, B.; Kartalov, E. and Quake, S.R.** (2003). Sequence information can be obtained from single DNA molecules. Proc. Natl. Acad. Sci. USA, 100, 3960–3964.
- Caetano-Anollés, G.** (1994). MAAP: a versatile and universal tool for genome analysis. Plant Mol. Biol. 25:1011-1026.
- Caetano-Anollés, G., B.J. Bassam, and P.M. Gresshoff** (1991). DNA amplification fingerprinting using very short arbitrary oligonucleotide primers. Bio/Technology 9:553-557.
- Dayrat, B.** (2005). towards integrative taxonomy. Biological Journal of the Linnean Society 85, 407–415.
- Demasure, B., Sodzi, N. and Petit, R.J.** (1995). A set of universal primers for amplification of polymorphic non-coding regions of mitochondrial and chloroplast D NA in plants. Molecular Ecology 4: 129-131.
- Deu, M., D. Gonzalez-de-Leon, J.C. Glaszmann, I. Degremont, J. Chantereau, C. Lanauad and P. Hamon.** (1994). RFLP diversity in cultivated sorghum in relation to racial differentiation. Theor. Appl. Genet. 88:838-844.
- Dong, J. and Wagner, D.B.** (1993). Taxonomic and population differentiation of mitochondrial diversity in *Pinus banksiana* and *Pinus contorta*. Theor. Appl. Genet. 86:573-578.
- Dunn, C.P.** (2003). Keeping taxonomy based in morphology. Trends Ecol Evol 18:270–271
- Hamby, R.K. and Zimmer, E.A.** (1992). Ribosomal RNA as a phylogenetic tool in plant systematics. In: Soltis PS, Soltis DE, Doyle JJ (eds) Molecular systematics of plants. Chapman & Hall, New York
- Hayashi, K.** (1992). PCR-SSCP: A method for detection of mutations. Genetic Analysis: Techniques and Applications 9:73-79.
- Hebert, P.D.N., Cywinska, A., Ball, S.L. and DeWaard, J.R.** (2003). Biological identifications through DNA barcodes. Proc R Soc B 270:313–321.
- Hillis, D.M. and Dixon, M.T.** (1991). Ribosomal DNA : molecular evolution and phylogenetic inference. Q Rev Biol 66:411–453
- Jack, P.L., Dimitrijevic, T.A.F. and Mayes, S.** (1995). Assessment of nuclear, mitochondrial and chloroplast RFLP markers in oil palm (*Elaeis guineensis* Jacq.). Theor. Appl. Genet. 90:643-649.
- Karvonen, P., Szmidt, A.E. and Savolainen, O.** (1994). Length variation in the internal transcribed spacers of ribosomal DNA in *Picea abies* and related species. Theor. Appl. Genet. 89:969-974.
- Lynch, M.** (1990). The similarity index and DNA fingerprinting. Mol. Biol. Evol. 7:478-484.
- Lynch, M. and Milligan, B.G.** (1994). Analysis of population structure with RAPD markers. Mol. Ecol. 3:91-99.

*Corresponding Author

- Milligan, B.G.** (1991). Chloroplast DNA diversity within and among populations of *Trifolium pratense*. *Current Genet.* 19:411-416.
- Moritz, C. and Cicero, C.** (2004). DNA barcoding: promise and pitfalls. *PLoS Biol* 2:1529–1531.
- Morgante, M. and Olivieri, A.M.** (1993). PCR-amplified microsatellites as markers in plant genetics. *Plant J.* 3:175-182.
- Poczai, P. and Hyvönen, J.** (2010). Nuclear ribosomal spacer regions in plant phylogenetics: problems and prospects. *Mol Biol Rep* 37:1897–1912.
- Qi, X., Stam, P. and Lindhout, P.** (1998). Use of locus specific AFLP markers to construct a high density molecular map in barley. *Theor Appl Genet* 96:376–384.
- Saal, B. and Wricke, G.** (2002). Clustering of amplified fragment length polymorphism markers in a linkage map of rye. *Plant Breed* 121:117–123.
- Saghai-Marcoof, M.A., Biyashev, R.M., Yang, G.P., Zhang, Q. and Allard, R.** (1994). Extraordinarily polymorphic microsatellite DNA in barley: Species diversity, chromosomal locations and population dynamics. *Proc. Nat. Acad. Sci. USA* 91:5466-5470.
- Saliba-Colombani, V., Causse, M., Gervais, L. and Philouze, J.** (2000). Efficiency of RFLP, RAPD, and AFLP markers for the construction of an intraspecific map of the tomato genome. *Genome* 43:29–40.
- Savolainen, V., Cowan, R.S., Vogler, A.P., Roderick, G.K. and Lane, R.** (2005). Towards writing the encyclopaedia of life: an introduction to DNA barcoding. *Philos Trans R Soc B Biol Sci* 360:1805–1811.
- Setoguchi, H.; Mitsui, Y.; Ikeda, H.; Nomura, N.; Tamura, A.** (2009). Development and characterization of microsatellite loci in the endangered *Tricyrtis ishiiana* (Convallariaceae), a local endemic plant in Japan. *Conserv. Genet.*, 10, 705–707.
- Shriner, M.O., Jin, L., Chakraborty, R. and Boerwinkle, E.** (1993). VNTR allele frequency distributions under the stepwise mutation model: a computer simulation approach. *Genetics* 134:983–993.
- Varshney, R.K.; Chabane, K.; Hendre, P.S.; Aggarwal, R.K. and Graner, A.** (2007). Comparative assessment of EST-SSR, EST-SNP and AFLP markers for evaluation of genetic diversity and conservation of genetic resources using wild, cultivated and elite barleys. *Plant Soc.* 2007, 173, 638–649.
- Welsh, J. and McClelland, M.** (1990). Fingerprinting genomes using PCR with arbitrary primers. *Nucleic Acids Res.* 18:6531-6535.
- Williams, J.G.K., Kubelik, A.R., Livak, K.J., Rafalski, J.A. and Tingey, S.V.** (1990). DNA polymorphism amplified by arbitrary primers are useful as genetic markers. *Nucleic Acid Res.* 18:6531-6535.
- Will, K.W. and Rubinoff, D.** (2004). Myth of the molecule: DNA barcodes for species cannot replace morphology for identification and classification. *Cladistics* 20:47–55.
- Young, W.P., Schuppet, J.M. and Keim, P.** (1999). DNA methylation and AFLP marker distribution in the soybean genome. *Theor Appl Genet* 99: 785–792.