ISOLATION AND MOLECULAR CHARACTERIZATION OF PLANT GROWTH PROMOTING RHIZOBACTERIA FROM THE HIGH ALTITUDE HIMALAYAN REGION OF UTTARAKHAND

Prashant Singh^{1, 7}, Deep Chandra Suyal², Naveen Kumar³, Manish Dev Sharma⁴, Indra Rautela⁵, Aditya Patel,⁶ Narottam Sharma⁷ and Veena Pandey¹

¹Department of Biotechnology, Bhimtal Campus, Kumaun University, Nainital, Uttarakhand ²Department of Microbiology, Govind Ballabh Pant University of Agriculture & Technology, Pantnagar, Uttarakhand.

³Molecular Cytogenetics Laboratory, Department of Molecular Biology & Genetic

Engineering, Govind Ballabh Pant University of Agriculture & Technology Pantnagar.

⁴Department of Life Sciences, Shri Guru Ram Rai Institute of technology & Science, Dehradun

⁵Uttaranchal College of Applied and Life Sciences, Uttaranchal University, Dehradun

⁶Devsthali Vidhyapeeth, Lalpur, U S Nagar

⁷Central Molecular Research Laboratory, Biochemistry Department, Shri Guru Ram Rai Institute of Medical & Health Sciences, Dehradun Email: Veena_kumaun@yahoo.co.in

Received-10.02.2017, Revised-02.03.2017

Abstract: The objective of this study was to isolate and characterize a rhizospheric bacterium from Munsyari, (2200 feet, 30.06°N/80.23° E) Uttarakhand, western Himalayas, (India). Plant growth promoting rhizobacteria (PGPR) are known to influence plant growth by various direct or indirect mechanisms. Isolated strain was tested for various PGP traits like 1-aminocyclopropane-1-carboxylate (ACC) deaminase activity, phosphate solubilisation, indole acetic acid production, production of siderophore, carbohydrate utilization test. Bio-control ability of isolate was also screened. Further identification of isolate was performed by PCR based 16S rRNA gene sequencing. The isolate PS03 was found to be most effective.

Keywords: Rhizobacteria, Isolation, Molecular, Himalaya region

REFERENCES

Dobbelaere, S.; Vanderleyden, J. and Okon, Y. (2003). Plant growth promoting effects of diazotrophs in the rhizosphere. Crit Rev Plant Sci 22:107–149 doi:10.1016/j.soilbio.2004.08.030

Dobbelaere, S.; Vanderleyden, J. and Okon, Y.(2003). Plant growth-promoting effects of diazotrophs in the rhizosphere. CRC Crit Rev Plant Sci.22:107–149

Edi-Premono Moawad, M. and Vleck, P.L.G. (1996). Effect of phosphate solubilizing *Pseudomonas putida* on the growth of maize and its survival in the rhizosphere. *Indonesian Journal of Crop Science*. 11:13–23.

Glick, B.R. (1995). The enhancement of plant growth by free living bacteria. Can J Microbiol 41:109–117

Gray, E.J. and Smith, D.L. (2005). Intracellular and extracellular PGPR: Commonalities and distinctions in the plant-bacterium signaling processes. Soil Biol Biochem. 37:395–412.

Gray, E.J. and Smith, D.L. (2005). Intracellular and extracellular PGPR:commonalities and distinctions in the plant-bacterium signaling processes. Soil BiolBiochem 37:395–412.

Greiner, R. and Alminger, M. L. (2001). "Stereospecificity of myo-inositol hexakisphosphatedephosphorylation by phytatedegrading enzymes of cereals," Journal of Food Biochemistry, 25 (3): 229–248'

Hayat, R., Ali, S., Amara, U., Khalid, R., Ahmed, I. (2010). Soil beneficial bacteria and their role in plant growth promotion: a review. Ann Microbiol 60:579–598. doi:10.1007/s00248-007-9247-9

Kerovuo, J., Lauraeus, M., Nurminen, P., Kalkkinen N. and Apajalahti, J. (1998). "Isolation, characterization, molecular gene cloning, and sequencing of a novel phytase from Bacillus subtilis," Applied and Environmental Microbiology, 64 (6) 2079–2085

Neumann, B., Pospiech, A. and Schairer, H.U. (1992). Rapid isolation of genomic DNA from gramnegative bacteria. Trends Genet TIG 8(10):332

Patten, C.L. and Glick, B.R. (2002). Role of Pseudomonas putidaindoleacetic acid in development of the host plant root system. *Applied and Environmental Microbiology*. 68(8):3795–3801.

Pikovskaya, R.I. (1948). Mobilization of phosphorus in soil in connection with vital activity of some microbial species. *Microbiology*. 17:362–370.

Richardson, A. E.; Hadobas, P. A.; Hayes, J. E.; O'hara, C. P. and Simpson, R. J. (2001). "Utilization of phosphorus by pasture plants supplied with myo-inositol hexaphosphate is enhanced by the presence of soil micro-organisms," Plant and Soil, 229(1) 47–56

*Corresponding Author

218 PRASHANT SINGH, DEEP CHANDRA SUYAL, NAVEEN KUMAR, MANISH DEV SHARMA, INDRA RAUTELA, ADITYA PATEL, NAROTTAM SHARMA AND VEENA PANDEY

Schwyn, B. and Neilands, J.B. (1987). Universal chemical assay for the detection and determination of siderophores. *AnalyticalBiochemistry*. 160(1):47–56.

United Nations, Department of Economic and Social Affairs, Population Division (2015). World Population Prospects: The 2015 Revision, Methodology of the United Nations Population Estimates and Projections. ESA/P/WP.242.

Van Loon, L.C. (2007). Plant responses to plant growth-promoting rhizobacteria. Eur J Plant Pathol 119:243–254. doi:10.1007/s10658-007-9165-1 Vessey, J.K. (2003). Plant growth promoting

rhizobacteria as biofertilizers. Plant Soil 255:571– 586. doi:10.1023/A:1026037216893