SURVEY AND SURVEILLANCE OF WHEAT POWDERY MILDEW IN MAJOR GROWING AREAS OF NORTHERN HILLS AND PLAINS, INDIA

Vipin Kumar Sharma*, A.S Rathi¹, M.S. Shahran², J. Singh¹ and S.S. Karwasra¹

¹Department of Plant Pathology, CCS Haryana Agricultural University, Hisar-125004, India ²Indian Institute of Wheat and Barley Research, Karnal-132001, India Email: mr.vipinsharma007@gmail.com

Received-02.08.2019, Revised-28.08.2019

Abstrct: Powdery mildew is very important foliar disease of wheat among major wheat growing areas of northen hills and plains in India which results in huge economic loss. Disease first appear on the lower leaves, varied in appearence depends on the location and continue to produce white mycelial growth on all plant parts up to the maturity of crop. Survey results revealed that the disease severity of powdery mildew in all the wheat growing areas of northern part of India is low to moderate (0 to 7). The disease severity was varied among different stage of the crop and places to grown but there was no significant difference among the cultivation practices. Maximum disease severity grade was recorded in Dhaulakuan district *i.e* 6 in 2012-13 and 7 in 2013-14, whereas minimum disease severity was recorded in Ludhiana, Ambala and Kaul i.e. 3 and 2 during the year 2012-13 and 2013-14, respectively.

Keywords: Powdery mildew, Northern hills, Survey, Wheat

REFERENCES

Anonymous(2018). Agriculture and food. <u>https://www.agric.wa.gov.au/spring/managing-</u>powdery-mildew-wheat.

Awad, Y.M., Abdullah, A.M., Bayoumi, T.Y., Elsalam, K. and Hassanien, A.E. (2015). Early Detection of Powdery Mildew Disease in Wheat (*Triticum aestivum* L.) Using Thermal Imaging Technique. Advances in Intelligent Systems and Computing. 323, DOI: 10.1007/978-3-319-11310-4 66.

McDonald, B.A. and Linde, C. (2002). Pathogen population genetics, evolutionary potential and durable resistance. Annual Review Phytopathology. 40:349–379.

Mehta, K.C. (1930). Studies on annual recurrence of powdery mildew of wheat and barley in India. Agricultural Research Journal. 25: 283-286

Naik, H.P. and Kulkarni, S. (2018). Survey of Powdery Mildew in Major Cucumber Growing Areas of Northern Karnataka, India. International Journal of Current Microbiology and Applied Sciences. 7(4): 3015-3024.

Rajak RL, Diwakar MC. (1995). National surveillance on crop pest and diseases.National seminar on changing pest situation in the current agriculture scenario of India. Ed. 1, Vol 1, ICAR, Pusa, New Delhi. 3-20.

Sankaran, S., Mishra, A., Ehsani, R. and Davis, C. (2010). A review of advanced techniques for detecting plant diseases. Computers and Electronics in Agriculture. 72:1–13.

Sharma, T.R., Singh, B.M. and Basandrai, A.K. (1992). Role of cleistothecia in annual recurrence of wheat powdery mildew in north India. Indian Phytopathology. 45(2): 205-206.

West, J.S., Bravo, C., Oberti, R., Lemaire, D. and Moshou, D. (2003). The potential of optical canopy measurement for targeted control of field crop diseases. Annual review of Phytopathology. 41:593– 614.

Zohary, D., Hopf, M. and Weiss, E. (2012). Domestication of Plants in the OldWorld: The origin and spread of domesticated plants in Southwest Asia, Europe, and the Mediterranean Basin. Oxford University Press.

Coakley, S.M., Scherm, H. and Chakraborty, S. (1999). Climate change and plant disease management. Annual Review of Phytopathology 37(1), 399–426.

Strange, R.N. and Scott, P.R. (2005). Plant disease: a threat to global food security. Phytopathology; 43

Braun, U., Cook, R., Inman, A., Shin, H., B'elanger, R., Bushnell, W., Dik, A. and Carver, T. et al. (2002). The taxonomy of the powdery mildew fungi. The powdery mildews: a comprehensive treatise, pp. 13–55.

Zhang, J.C., Pu, R.L., Wang, J.H., Huang, W.J., Yuan, L. and Luo, J.H. (2012). Detecting powdery mildew of winter wheat using leaf level hyperspectral measurements. Computers and Electronics in Agriculture 85, 13–23.

Cooke, B.M., Jones, D.G. and Kaye, B. (2006). The Epidemiology of Plant Diseases. NetherlandsSpringer 57. Coops NC, Wulder MA, White JC (2006) Integrating remotely sensed and ancillary data sources to characterize a mountain pine beetle infestation. Remote Sensing of Environment, 105: 83–97.

Mehta, N., Singh, K. and Sangwan, M. S. (2009). Resistance to powdery mildew of mustard in relation to rate reducing factors. *Plant Disease Research*.24(2): 114-119.

*Corresponding Author

Journal of Plant Development Sciences Vol. 11(8): 467-470. 2019