KNOWLEDGE AND ADOPTION OF RECOMMENDED MAIZE PRODUCTION TECHNOLOGY

P.K. Netam*, H.K. Awasthi and R.S. Sengar

1,2,3,Department of Agricultural Extension CoA, IGKV, Raipur, Chhattisgarh Email: pknetam49@gmail.com

Received-06.12.2018, Revised-25.12.2018

Abstracts: This investigation was carried out in three district of Bastar plateau of Chhattisgarh State to assess the level of knowledge and adoption of recommended maize production technology. 270 farmers were considered as respondents for this study. Respondents were interviewed through personal interview. Collected data were analyzed with the help of suitable statistical methods. The analysis of the results showed that overall knowledge of recommended maize production technology, 72.96% respondents had medium level of knowledge and 73.70% respondents had medium level of adoption regarding recommended maize production technology.

Keywords: Maize production, Knowledge, Adoption, Technology

REFERENCES

Bawa, D.B. and Ani, A.O. (2014). Analysis of Adoption of Improved Maize Production Technology among Farmers in Southern Borno, Nigeria. Research on Humanities and Social Sciences, 4(25): 137-141.

Chhattisgarh (2014). Annual statistics report.

CIMMYT (2005). Maize in India: production systems, constraints, and research priorities.

Gecho, Yishak and Punjabi, N.K. (2011). Determination of adoption of improved Maize technology in Damot Gale, Wolaita, Ethiopia. Raj. J. Ext. Edu., 19: 1-9.

Gupta, Km. Saroj and Gyanpur, S.R.N. (2012). Sustainability of scientific maize cultivation practice in Uttar Pradesh, India. Journal of Agricultural Technology. 8 (3): 1089-1098.

Langade, D. M., Shahi, J.P., Agrawal, V. K. and Sharma, A. (2013). Maize as emerging source of oil in india: an overview. Maydica, 58(3/4): 224-230.

Paikra, V. K. (2014). Assessment of technological gap in production of black gram among the tribal farmers of Jashpur District Chhattisgarh. M. Sc. (Ag.) Thesis, IGKV, Raipur.

R. Cox, (1956). Control of helminthosporium turcicum blight disease of sweet corn in South Florida. Phytopathology, 5: 68-70.

Reddy, T. R., Reddy, P. N., Reddy, R. R. and Reddy, S. S. (2013). Management of Turcicum leaf blight of maize, caused by Exserohilum Turcicum in maize. International Journal of scientific and Research publications, 3(10): 1-4.

Willy, V. (Undated). Soil plant growth and production Vo. II National Science foundation Flanders and geography department, Belgium: University of Ghenl . (accessed on 02/01/2013).

Yadav, S., Prajapati, R. R. and Prajapati, M.R. (2014). Knowledge and adoption of tomato growers about improved tomato production technology. Guj. J. Ext. Edu., 25(2): 172-174.

*Corresponding Author