PHYSICAL CHARACTERISTICS AND CHEMICAL CONSTITUENTS IN CRUDE PLANTS METHANOLIC EXTRACT OF SELECTED MEDICINAL PLANTS BASED ON LITERATURE AND TRADITIONAL KNOWLEDGE

V.P.S. Bhadauria*, and Varsha Gupta¹

Department of Chemistry, St. John's College, Agra-282002, India ¹Agronomy, RVSKVV, Gwalior Chapter at Foundation for Innovative Research, Sustainable Technologies and Intellectual Property (FIRSTIP) www.firstip.org Email: <u>vps.chem@gmail.com</u>

Received-06.04.2019, Revised-26.04.2019

Abstract: Eucalyptus globules yielded highest in case of leaves extract, while in case of fruit extract Annona squamosa yield highest and in case of seed extract Butea frondosa yield was highest. Phytochemical evaluation showed that Ailenthus excels (leaves) extract was positive for alkaloids, flavonoids, tannins, phenolic compounds and triterpenoids, while in Calotropis procera (leaves) methanolic extract alkaloids, flavonoids and tannins were identified and in case of Chenopodium album (seeds) alkaloids, saponins, glycosides, fixed oils and tannins were present.

Keywords: Phytochemical, Medicinal plants, Alkaloids, Flavonoids, Tannins, Phenolic compounds, Triterpenoids

REFERENCES

Chung, A.C. (1995). Antimicrobial Agents. Chemother., **39**: 2235-2238

Chopra, R.N. and Chopra, I.C. (1955). A review of work on Indian medicinal plants. I.C.M.R. special reports series, pp 102

Mc. Corkle, C.M. (1982). Management of animal health and disease in an indigenous anderson community SRC RSP Technical report No. 4 Department of Rural Sociology, University of Missouri, Columbia, Missouri, USA pp 106.

Misri, J., Vihan, V.S., Kumar, A. and Sharma, D.K. (2002). Evaluation of anthelmintic properties of *M. azedarach* (Bakain) on experimental *H. contortus* infection in goats. Indian J. Vet. Med., **22** (2), 4-6

Malairajan, P., Gopalakrishnan., Geetha., Narasimhan, S., Jessi., Kala and Ven. K. (2006). Analgesic activity of some Indian medicinal plants. J. Ethnopharmacol., **106(3):** 425-428.

Kosalge, S.B. and Fursule, R.A. (2009). Investigation of ethnomedicinal claims of some

plants used by Tribals of Satpuda Hills in India. J. Ethnopharmacol., **121(3):** 456-461.

Debela, A. (2002). Mannual of Phytochemical screening of medicinal plants. Ethiopian Health and Nutrition Research Institute, Addis Ababa, Ethiopia, 35-47

Paech, K. and Tracy, M.V. (1955). Modern methods of plant analysis. III.

Thanabhorn, S., Jenjoy, K., Thamaree, S., Ingkaninan, K. and Panthong, A. (2006). Acute and subacute toxicity study of the ethanol extract from *Lonicera japonica*. J. Ethnopharmacol., **107**: 370-373

Rajesh, M.G. and Latha, M.S. (2004). Preliminary evaluation of the antihepatotoxic activity of Kamilari, a ployherbal formulation. J. Ethnopharmacol., **91**: 99-104.

Gupta, A.K. and Mishra, N. (2006). Hepatoprotective activity of aqueous ethanolic extract of *Chamomile capitula* in paracetamol intoxicated albino rats. Am. J. Pharmacol. Toxicol., **1**: 17-20.

*Corresponding Author

Journal of Plant Development Sciences Vol. 11(4): 253-256. 2019