

**SURVEY ON POWDERY MILDEW (*ERYSIPHE POLYGONI* DC) DISEASE IN
CORIANDER (*CORIANDRUM SATIVUM* L.) AT GWALIOR DIVISION**

Rajendra Kashyap¹, P.K. Bhagat² and G.P. Painkra^{2*}

¹*Indira Gandhi Krishi Vishwavidyalaya, Krishi Vigyan Kendra, Balrampur, Chhattisgarh, India*

²*Indira Gandhi Krishi Vishwavidyalaya, Rajmohini Devi College of Agriculture and Research
Station, Ambikapur, Distt- Surguja (C.G.) India 497001*

Received-22.09.2015, Revised-29.09.2015

Abstract: A survey on powdery mildew disease in coriander at Gwalior Division of Madhya Pradesh to assess the intensity of powdery mildew on the farmer's fields. It is an important disease of coriander in Guna district. Among the surveyed villages the minimum intensity of powdery mildew 14.73 per cent was recorded in Magroda village of Raghogarh block. However it was maximum 44.14 per cent in Bhadodi village of Raghogarh block. Out of the five Surveyed blocks the minimum disease intensity was recorded in bamori 20.79 per cent followed by Kumbhraj 27.22 per cent, Aron 30.18 per cent and Chachoda 34.79 per cent, while maximum intensity 36.65 per cent was recorded in Raghogarh block.

Keywords: *Coriandrum sativum*, Disease, Powdery mildew

REFERENCES

Das, S.R. and Narain, A. (1990). Management of powdery mildew of mung bean with fungicide. *Ind. Phytopath.* 43(1): 100-101

E1-Meleigi, M.A. and A1-Rokibah, A. (1996). Survey of Wheat Disease in Central Saudi Arabia. *Bulletin Facul. Agril. Univ Cairo* 47(3): 499-512.

Lo, C.T. and Wang, K.M. (2000). Survey of fungal diseases on Above ground parts of Wasabi in Taiwan. *Pl. Path. Bull.* 9(1):17-22.