

EVALUATION OF FUNGI TOXICANTS AGAINST POWDERY (*ERYSIPHE POLYGONI* DC) DISEASE IN CORIANDER (*CORIANDRUM SATIVUM* L.) AT GWALIOR DIVISION

Rajendra Kashyap¹, P.K. Bhagat² and G.P. Painkra^{2*}

¹. Indira Gandhi Krishi Vishwavidyalaya, Krishi Vigyan Kendra, Balrampur, Chhattisgarh, India

². Indira Gandhi Krishi Vishwavidyalaya, Rajmohini Devi College of Agriculture and Research Station, Ambikapur, Distt- Surguja (C.G.) India 497001

Email: gppainkrarmd@gmail.com

Received-20.09.2015, Revised-27.09.2015

Abstract: Five fungicides viz. wettable sulphar, carbendazim, difenoconazole, mancozeb and saaf were tested against powdery mildew (*E. poligoni*) of coriander. The minimum intensity of the disease was recorded in the treatment sulfex (19.86 per cent) followed by score 25.83 per cent, bavistin 29.72 per cent, saaf 31.67 per cent and mancozeb 39.30 per cent.

Keywords: Coriander, Fungi toxicants, Powdery mildew

REFERENCES

Keshwal, R.L., Choubey, D.C. and Singh, K. (1979). Effect of date of sowing and fungicidal sprays on powdery mildew of coriander. *Pesticides*. **6** (2) : 135-136.

Raju, K.S., Rao, M. G., Rao, T. S. and Babu, M. K. (1982). Note of fungicidal control of coriander powdery mildew. *Ind. J. Agric. Sci.* **52** (4) : 262-263.
Srivastava, U.S., Rai, R. A. and Agrawal, J. M. (1971). Powdery mildew of coriander and its control. *Indian Phytopath.* **24** (3) : 437-440.

*Corresponding Author