INTEGRATED WEED MANAGEMENT IN SYSTEM OF RICE INTENSIFICATION (SRI) FOR SECURED LIVELIHOOD

Devendra Kumar Dewangan¹, Shivam Soni², Harish Kumar Netam³

Department of Agronomy¹
Department of Genetics and Plant Breeding²
Department of Entomology³
Indira Gandhi Krishi Vishwavidyalaya, Raipur-492012, Chhattisgarh, India

Abstracts: Weeds not only cause quantitative, but also hamper the quality of produce owing to competition for nutrients, moisture, light and to some extent for space. The extent of yield reduction of rice due to weeds is estimated from 15-95 per cent (Gogai *et al.*, 1996).

Keywords: Livelihood, Rice, Weed

REFERENCES

Gogoi, A. K., Brown, H., Cussans, G. W., Devine, M. D., Duke, S.O., Fernandes, Q.C., Helweg, A., Labrada, R. E., Lanndes, M., Kudsk, P. and Steibig, J.C. (1996). Integrated weed management of rice in high rainfall region of India: Status and Prospects. *In*: Proceedings of the second International Weed Control Congress, Copenhagen, Denmark, 25-28 June. 1-4: 715-719.

Kolhe, S.S. (1999). Evaluation of low dosage-high efficacy herbicides Fenoxaprop-p-ethyl and ethoxysulfuron in direct seeded rice under puddle condition. *Oryza.* **36** (2):177-179.

Moorthy, B. T. S. Saha S. (2002). Bio-efficacy of Certain New Herbicide Formulations in Puddle-seeded Rice. *Indian Journal of Weed Science*: **34**(1-2)