ANALYSIS OF QUALITATIVE TRAITS IN OKRA [ABELMOSCHUS ESCULENTUS (L.) MOENCH] GROWN UNDER TWO ENVIRONMENTS

Sankara Rao Karri and Pinaki Acharyya

Department of Horticulture, Institute of Agricultural Science University of Calcutta, 35. Ballygunge Circular Road, Kolkata – 700019, India E-mail: ksrgreenplus@hotmail.com, pinakiacharyya@yahoo.co.in

Abstract: Besides Okra is a potential fibre yielding crop as because the bast fibre is strong, hydroscopic and resistant to rot, thus suitable to meet the global demand as an additional source of ecofriendly fibre. Fifteen genotypes of Okra were evaluated for morphological and yield related traits. Estimation of biochemical constituents i.e. total soluble solids, crude fibre, total cartenoids, calcium and phosphorus were also performed.

Keywords: Abelmoschus esculentus, Analysis, Okara

REFERENCES

Association of Official Agricultural Chemists (1975). Officer and Tentative Methods of Analysis.

Chapman,H.D.,Pratt.,P.F.,(1961). Methods of analysis for Soils, *plants and water univ*.California, Berkeley,CA.,USA

Joshi, A.B. and Hardas, M.W. (1976). Okra. In : *Evolution of crop plants* (Ed. N.W. Simmonds), Longmans, London, P. 194.

Mahlberg, P.G. and Venkateswarm, S. (1966). Pigment analysis of normal and proliferated genetical strains of Nicotiana under cultural conditions. *Bot. Gaz.*, **127**: 114 - 119.

Patil, Y.B., Madalageri, B.B., Biradar, B.D. and Hosamani, R.M. (1996). Variability Studies in Okra [*Abelmoschus esculentus* (L.) Moench]. *Kar. J. Agric. Sci.*, 9 (2) : 289 – 293.

Rao, T.S. (1972). Note on the Natural Variability for some Qualitative and Quantitative Characters in Okra. *Ind. J. Agri. Sci.*, **42** : 437 – 438.

Sood, S., Ariya, P.S. and Singh, Y. (1995). Genetic Variability and Correlation Studies in Okra. [*Abelmoschus esculentus* (L.) Moench]. *Advance in Horticulture and Forestry.*, **4** : 109 – 118.