EFFECT OF COPPER AND MERCURY ON GROWTH PARAMETERS IN LEMNA MINOR

Smita Pathak and Abha Agarwal

Department of Botany, Bareilly College, Bareilly (U.P.) Smita_111pathak@yahoo.com

Abstract : Theability of aquatic plants to accumulate heavy metals from water is well documented. In this study, duckweeds plants (*Lemna minor*) were exposed to different concentrations of Cu and Hg. Various growth parameters (fresh weight, dry weight and growth index) in different seasons(summer and winter) were studied. The effect of all concentrations on plant parameters was toxic. The plant growth was decreased as the concentrations of heavy metals were increased. The results suggest that the *L.minor* can be effectively used as phytoremediator for waste water polluted with more than one heavy metal at moderate concentrations.

Keywords : Cu and Hg, Heavy metal, Lemna minor, Plant-toxicity, Phytoremediation

REFERENCES

An Y.J., Kim Y.M., Kwon T.I. and Jeong S.W. (2004). Combined effects of copper, cadmium and lead upon *Cucumis sativus* growth and bioaccumulation. *Sci.Total Environ.*, **326**:85-93.

Balsbery Pahlson A.M.(1989). Toxicity of heavy metals (Zn,Cu,Cd,Pb) to vascular plants. *Water Air Soil Pollut.*,**47**:287-319.

Cheng J., Landesman L., Bergmann B.A., Classen J.J., Howard J.W. and Yamamoto Y.T. (2002). Nutrient removal from swine lagoon liquid by *Lemna minor* 8627. *Transactions of the ASAE*, 45: 1003-1010.

Chiba S. and Takashasi, K.(1977). Studies on heavy metals pollution in agricultural land. (2) Absorption of cadmium and growth retardation in forage crops. *Bull. Shikoku.Agri. Exp. Station.*30:49-73.

Classen J.J., Cheng J., Bergmann B.A. and Stomp A.M. (2000).*Lemna gibba* growth and nutrient uptake in response to different nutrient levels. In: Animal, Agriculturaal and Food Processing Waste: Proc. Eighth International Symposium.pp:40-45. Des. Moines, Iowa. St. Joseph, Mich, ASAE.

Dirilgen W.(2011). Mercury and Lead : assessing the toxic effects on growth and metal accumulation by *Lemna minor*. *Ecotoxicol Environment Safety*,**74**(1) : 48.54.

Fridovich I. (1986). Biological effects of the superoxide radical. Archives of Biochemistry and Biophysics, 24:1-11.

Gardea – TorresdeyJ.I., Peralta- Videa J.R., Rosa G.D and Parsons J.G. (2005). Phytoremediation of heavy metalsand study of the metal coordination of heavy metals and study of the metal coordination of X-ray absorption spectroscopy. *Coord. Chem.*. Rev., 249 (17-18): 1797-1810.

Haele H.L. and Ormrod D.P. (1983). Effects of nickel and copper on seed germination, growth and development seedlings of *Acer pinnata*, *Betula papyrifera Picea abies and Pinus banksiana*. *Reclamation and Revogetation Research*, **2**: 41-54.

Kabata- Pendias A. and Pendias H. (2001). Trace elements in soil and plants. Boca Raton, FL : CRC press.

Khellaf N. and Zerdaoui M. (2009). Growth response of the duckweed Lemna minor to heavy metal pollution. *Iran J. Environ. Health Sci. Engineering*, 6(3):161-166.

Landesman L. (2000). Effects of herbivory and competition on growth of *Lemna*in systems for Wastewater Treatment and Livestock Feed Production, University of Louisiana at Lafayete, Louisiana, 150 pp

Lipman C.B. and Mackinney G. (1931). Proof of the essential nature of copper for higher green plants. *Plant Pysiol*, **6**: 593-599.

Lablora S., Mucha A. and Sprihoulova E. (1986). Compartmentation of cadmium, lead, zinc in seedlings of maize (*Zeamays* L.) and induction of metallothionin. *Biol. Czechas*, **4**:77-88.

Mkandawire M. (2005), Fate and Effect of Uranium and Arsenic in surface Waters of Abandoned Uranium Mines : Studies with *Lemna gibba* L. *Shaker, Aachem*, 185 pp.

Mkandawire M. and Dudel E. G. (2007). Are *Lemna* spp. Effective Phytoremediation Agents ? *Bioremediation, Biodiversity and Bioavailability*, 56-71.

Miretzky P., Saralegui A. and Cirelli A.F.(2004). Aquatic macrophytes potential for the simultaneous removal of heavy metals (Buenus Aires, Argentina), *Chemosphere*, **57**:997-1005.

Mkandawire M., Brackhage C., Taubert B. and Dudel E.G. (2005) . Semicontinuous culture system for *Lemna gibba* bioassay : functioning and theory of operation. *Applied Ecology and Environmental Research*, **3**:19-27.

Peralta –Videa J.R., Gardea- Torresdey J.L., Gomez E., Tiermann K.J., Parson J.G, and CarrilloG. (2002). Effect of mixed cadmium, copper, nickel and zinc at different pH upon alfalfa growth and heavy metal uptake. *Environ. Pollut.*, 119:291-301. **Quzounidou G., Elftheeriou, E. and Karaglis, S.** (1992) Ecophysiological and ultrastructural effects of copper in Thlaspi Ochroleucum (Cruciferal), *Can. J. Bot.*, **70:**947-957.

Rahman G.N.H. and Sternberg S.P.K. (1999). Bioremoval of lead using Lemna minor.*Bioresour. Technol.*, **70**:225-230.

Servillia Oancea, N. Foca and A. Airinei(2005). Effects of heavy metals on plant growth and photosynthetic activity.*Tomul I,s. Biofizica, Fizica medicala siFizica mediului*. pp 107-110.

Somner A.L. (1931).Copper as an essential metal for plant growth. *Plant physiology*, **6**:339-345.

Suszeynsky E.M. and Shann J.R. (1945). Phytotoxicity and accumulation of mercury in tobacco subjected to different exposure routes. *Environmental toxicology and chemistry.*,**14**:61-67.

Ting Y.P., Lawson F. and Prince I.G. (1991). Uptake of cadmium and zinc by algae *Chlorella vulgaris* : Multiion situation. *Biotechnical Bioeng.*, 37:445 -455.

Teisseire H. and Vernet G. (2000). Is the "diuron effect " due to a herbicide strengthening of antioxidative defense of *Lemna minor? Pesticide Biochem. Physiol.*, **66**:163-160.

Vyas J. and Puranik R.M. (1993). Inhibition of nitrate reductase activity by mercury in bean leaf segments. *Indian Journal of Plant Physiology*,**36**:57-60.

Waisberg M., Joseph P., Hale B. and Beyersmann D. (2003). Molecular and cellular mechanisms of cadmium carcinogenesis. *Toxicology*, **192**:95-117.

Zayed A., Growthaman S. and Terry N. (1998). Phytoaccumulation of trace elements by wetland plants : I. Duckweed. *J. Environ. Qual.*, **27**:715-721.