ROLE OF TURFGRASS IN URBAN LANDSCAPES

Sithin Mathew*

ICAR – Directorate of Floricultural Research, Pune – 411005 Email: sithin.mathew.m@gmail.com

Received-28.03.2021, Revised-02.05.2021, Accepted-22.05.2021

Abstract: Lawns occupy a significant proportion of green spaces in manycities worldwide. Despite of aesthetic and ornamental benefits, turfgrass also possess many inevitable functional benefits. It helps in maintaining ecological balance and reclamation of polluted environments due to profuse exploitation of nature in urban landscape. The major benefits include controlling soil erosion and improving soil quality, water purification and noise reduction, mitigating air pollution and dust. Turfgrass plays a major role in carbon sequestration which reduce the atmospheric temperature. Apart from these functional benefits, turfgrass is also used in various medical therapies as a tool for reducing mental stress and by various entrepreneurs as a billion-dollar industry with high returns per unit area.

Keywords: Turfgrass, Urban landscape, Air pollution, Water quality

REFERENCES

Alexander, M. (1997). Introduction to soil microbiology, 2nd Edn. John Wiley and Sons, New York. pp: 12-18.

Alireza, S. and Mohammad, H. (2019). The Effect of Rapid Population Growth on Urban Expansion and Destruction of Green Space in Tehran from 1972 to 2017. Journal of the Indian Society of Remote Sensing. https://doi.org/10.1007/s12524-019-00966-y.

Angle, J.S. (1985). Effect of cropping practices on sediment and nutrient loses from tobacco. Tobacco Science, 29: 107-110.

Artina, S., Bardasi, G., Borea, F., Franco, C., Maglionico, M. and Paoletti, A. (1999). Water quality modelling in ephemeral streams receiving urban overflows. In: The Pilot Study in Bologna. Proc. 8th International Conference on Urban Storm Drainage, ICUSD, Sydney. 1589–1596.

Athanasios, A.G., Cristiana, M.C., Mihai, R.N., Diana, A.O. and Irina, I.N. (2016). Urban Landscape Quality Index – planning tool for evaluating urban landscapes and improving the quality of life. Procedia Environmental Sciences, 32: 155-167.

Beard, J.B. and Johns, D. (1985). The comparative heat dissipation from three typical urban surfaces: Asphalt, concrete and a bermudagrass turf., In Texas Turf grass Research, Texas agricultural experiment station, pp-4329, college station.

Beard, J.B. (1973). Turfgrass: Science and Culture. Prentice-Hall, Englewood cliffs, NJ.

Bennett, H.H. (1939). Soil conservation, McGraw-Hill, Book, Co., New York.

Brady, N.C. and Weil, R.R. (1999). The Nature and Properties of Soils, 12th edition. Prentice Hall, New Jersey, USA. Pp. 100-108.

Cai, W., Liu, C. and Zhang, C. (2018). Developing the ecological compensation criterion of industrial

solid waste based on emergy for sustainable development. Energy, 157:940-948.

Chen, G., Wang, Y. and Li, S. (2018). Spatiotemporal patterns of PM 10 concentrations over China during 2005-2016: a satellite-based estimation using the random forests approach. Environmental Pollution, 242: 605.

Cheng, Z., Richmond, D.S., Salminen, S.O. and Grewa, P.S. (2008). Ecology of urban lawns under three common management programs. Urban Ecosystems, 11: 177–195.

Cook, D.I. and Van, D.F. Haverbeke, (1971). Trees and shrubs for noise abatement. Nebraska Agric. Exp. Stn. Res. Bull. 246, Lincoln.

Currie, B.A. and Bass, B. (2008). Estimates of air pollution mitigation with green plants and green roofs using the UFORE model. Urban ecosystem, 11(4): 409-422.

Dabney, S.M., Delgado, J.A. and Reeves, D.W. (2001). Using winter cover crops to improve soil and water quality. Communications in Soil Science and Plant Analysis, 32: 1221-1250.

Ehleringer, J.R. and Cerling, T.E. (2002). C3 and C4 photosynthesis In Encyclopedia of Global Environmental Change. The Earth System: Biological and Ecological Dimensions of Global Environmental Change, R. E. Munn, eds., Wiley, New York, pp. 186–190.

European Council, European landscape convention, E. Council, Editor (2000). European Council

Fontaine, S., Mariotti, A. and Abbadie, L. (2003). The priming effect of organic matter: a question of microbial competition. Soil Biology and Biochemistry, 35(6): 837-843.

Ghannoum, O., Caemmerer, S.V., Ziska, L.H. and Conroy, J.P. (2000). The growth response of C4 plants to rising atmospheric CO₂ partial pressure: a reassessment. Plant Cell Environ, 23(9): 931–942.

Gould, F.W. (1968). Grass systematics, McGraw Hill, New York.

*Corresponding Author

- **Gross, C.M., Angle, J.S. and Welterlen, M.S.** (1990). Nutrient and sediment losses from turf grass. Journal of Environmental Quality, 19: 663-668.
- **Gross, C.M., Angle, J.S. and Welterlen, M.S.** (1991). Run off and sediment losses from tall fescue under simulated rainfall. Journal of Environmental Quality, 20: 604-607.
- **Gu, C., Cranell, J., Hornberger, G. and Carrico, A.** (2015). The effects of household management practices on the global warming potential of urban lawns. Journal of Environmental Management, 151: 233-242.
- **Jain, V.K.** (1983). Plant physiology. S. Chand and Company, New Delhi.
- **James, B.B. and Robert, L.G.** (1994). The Role of Turfgrasses in Environmental Protection and Their Benefits to Human. Journal of Environmental Quality, 23:452-460.
- **Janakiram, T., Ritu, J. and Namita.** (2015). Introduction to turf grasses. Westville, New Delhi, pp. 12-15.
- **John, C.** (2013). World Population Growth, Past, Present and Future. Environ Resource Economy, 55:543-554
- **Johns, D. and Beard, J.B.** (1985). Aquantitative assessment of the benefits from irrigated turf on environmental cooling and energy savings in urban areas. P. 134-142. In Texas Turf grass Research, Texas agricultural experiment station, Pr-4330, college station.
- **Johnson. J. and Newton, J.** (1996). Building green, a guide for using plants on roofs walls and pavement, the London ecology unit, London.
- **Jose, A.M.** (2017). Ecosystem services from turfgrass landscapes. Urban forestry and Urban Greening, 16: 30139.
- **Keeley, E. and Rundel, P.W.** (2005). Fire and the Miocene expansion of C4 grasslands. Ecological Letters, 8(7): 683–690.
- Kim, R.H., Lee, S., Kim, Y.M., Lee, J.H., Kim, S.K. and Kim, S.G. (2005). Pollutants in rainwater runoff in Korea: their impacts on rainwater utilization. Environmental Technology, 26: 411-420.
- **Lai, L., Huang, X. and Yang, H.** (2016). Carbon emissions from land-use change and management in China between 1990 and 2010. Science Advances, 2 (11).
- **Lamb, H.H.** (2010). Industrial smoke drift and weather. Quarterly Journal of Royal Meteorological Society, 64(277):639-643.
- **Lee, J.H. and Bang, M.K.W.** (2000). Characterization of urban storm water runoff. Water Research, 34(6): 1773–1780.
- **Lee, K.E.** (1985). Earthworms. Their relationships with ecology and soil and land use. Academic Press, New south wales, Australia.
- **Linlin, G., Liangyun, L., Dailiang, P., Yong, H., Quanjun, J. and Lingling, L.** (2012). Monitoring the distribution of C3 and C4 grasses in temperate grassland in northern China using moderate

- resolution imaging spectro radiometer normalized difference vegetation index trajectories. Journal of Applied Remote Sensing, 6: 112-123.
- **Liu, Q. and Wang, Q.** (2017). How China achieved its 11th Five-Year Plan emissions reduction target: a structural decomposition analysis of industrial SO_2 and chemical oxygen demand. Sci. Total Environment, 574: 1104-1116.
- **Longwu, L., Zhenbo, Wang. and Jiaxin, L.** (2019). The effect of urbanization on environmental pollution in rapidly developing urban agglomerations. Journal of Cleaner Production, 237: 117649.
- **Lush, W.M.** (1990). Turf growth and performance evaluation based on turf biomass and tiller density. Agronomy Journal, 82: 505-511.
- Manish, S., Joseph, Y., Lindsey, S., Veronica, A. and Cade, C. (2021). Soil microbial biomass and composition from urban landscapes in a semiarid climate. Applied Soil Ecology, 158: 103810.
- Maria, I., Fredrik, E., Tuula, E., Per, B. and Marcus, H. (2017). The lawn as a social and cultural phenomenon in Sweden. Urban Forestry & Urban Greening, 21: 213–223.
- **Michael, E.D. and John, C.C.** (2005). A field evaluation of rain garden flow and pollutant treatment. Water, Air, and Soil Pollution, 167: 123–138.
- Milesi, C., Running, S.W., Elvidge, C.D., Dietz, J.B., Tuttle, B.T. and Nemani, R.R. (2005). Mapping and modeling the biogeochemical cycling of turf grasses in the United States. Environmental Management, 36: 426–438.
- Minke, G. and Witter, G. (1982). Haeusermit Gruenem Pelz, Ein Hand buchzur Hausbegruenung, Verlag Dieter Fricke GmbH, Frankfurt (partially translated in English).
- Morton, T.J., Gold, A.J. and Sullivan W.M. (1988). Influence of over watering and fertilization on nitrogen losses from home lawns. Journal of Environmental Quality, 17: 124-130.
- **Nedved, M. and Jansz, J.** (2006). Waste water pollution control in the Australian mining industry. Journal of Clean Production, 14(12&13):1118-1120.
- **Ng, B.J.H., Zhou, J. and Giannis, A.** (2014). Environmental life cycle assessment of different domestic wastewater streams: policy effectiveness in a tropical urban environment. Journal of Environmental Management, 140:60-68.
- Nowak, D.J., McHale, P.J., Ibarra, M., Crane, D., Stevens, J. and Luley, C. (1998). Modeling the effects of urban vegetation on air pollution. In: Gryning, S.E., Chaumerliac, N. (Eds.), Air Pollution Modeling and its Application XII. Plenum Press, New York, pp: 399–407.
- **Parton, W.J., Stewart, J.W.B. and Cole, C.V.** (1988). Dynamics of C, N, P, and S in grassland soils: a model. Biogeochem, 5:109-131.
- **Potter, D.A., Bridges, B.L. and Gordon, F.C.** (1985). Effect of nitrogen fertilization on earth worm

- and microarthropod populations in Kentucky blue grass turf. Agronomy Journal, 77: 367-372.
- **Robinette, G.O.** (1970). Plants people and environmental quality. U.S Dep. Interior, National Park Service and Am. Soc. Land. Archit. Foundation, Washington, DC.
- Roshini, A., Chawla, S.L. and Sudha, P. (2017). Evaluation of warm season turf grasses for various qualitative and quantitative traits under Gujarat agroclimatic conditions. Indian Journalof Agricultural Sciences, 87:83-91.
- Schulze, E.D., Ellis, R., Schulze, W., Trimborn, P. and Ziegler, H. (1996). Diversity, metabolic types and δ13C carbon isotope ratios in the grass flora of Namibia in relation to growth form, precipitation and habitat conditions. Population Ecology, 106:352-369. Schuyler, T. (1987). Controlling urban runoff: A practical manual for planning and designing urban BMPs. Metropolitan Washington Council of Governments, Washington, DC.
- **Sharma, B.K., Solanki, G.S. and Chalise, M.K.** (2014). Carbon sequestration in a community managed forest of Chitwan national park's buffer zone., Biological Journal, 9(1): 46-54.
- **Shi, W., Yao, H. and Bowman, D.** (2006). Soil microbial biomass, activity and nitrogen transformations in a turfgrass chronosequence. Soil Biology and Biochemistry, 38(2): 311-319.
- Sithin, M., Seetharamu, G.K., Dileepkumar, M. and Satish, D. (2021). Grasses for sports grounds and its influence on playing quality: A review. Journal of Pharmacognosy and Phytochemistry, 10(2): 17-26.
- Sithin, M., Seetharamu, G.K., Patil, B.C., Dileep Kumar, M., Amruta, S.B., Satish, D. and Mukund, S. (2020b). Economic Analysis of Profitability in Turf Grasses for Different Methods of Planting. International Journal of Current Microbiology and Applied Sciences, 9(12): 2551-2554.
- **Sithin, M., Seetharamu, G.K., Satish, D., Dileep Kumar, M. and Mukund, S.** (2020a). Effect of weed management and methods of planting on weed population and establishment of turf grasses. Crop Research, 55(5 & 6): 250-261.
- Stewart, G.H., Ignatieva, M.E., Meurk, C.D., Buckley, H., Horne, B. and Brad dick, T. (2009).

- URban Biotopes of Aotearoa New Zealand (URBANZ) (I): Composition and diversity of temperate urban lawns in Christchurch. Urban Ecosystems, 12(3): 233-248.
- **Thompson, G.L. and Kao-Kniffin, J.** (2019). Urban grass land management implications for soil C and N dynamics: a microbial perspective. Frontiersin Ecology and Evolution, 7(315):1-15.
- **UN Population Division** (2011a). World population prospects: the 2010 revision. New York.
- **Vogel, J.C., Fuls, A. and Danin, A.** (1986). Geographical and environmental distribution of C3 and C4 grasses in the Sinai, Negev, and Judean deserts. Bio Control, 70: 258-265.
- Wang, Y., Tu, C., Li, C., Tredway, L., Lee, D., Snell, M., Zhang, X. and Hu, S. (2014). Turfgrass management duration and intensities influence soil microbial dynamics and carbon sequestration. International Journal of Agriculture Biology, 16: 139-145.
- **Wang, Z.B. and Fang, C.L.** (2016). Spatial-temporal characteristics and determinants of PM 2.5 in the bohai rim urban agglomeration. Chemosphere, 148: 148-162.
- Watschke, T.L. and Mumma, R.O. (1989). The effects of nutrients and pesticides applied to turf on the quality of runoff and percolating water. Pennsylvania state university., Environmental Resources Research Institute., ER. 8904., University park.
- **Wu, J.G., Xiang, W.N. and Zhao, J.Z.** (2014). Urban ecology in China: historical de-velopments and future directions. Landscape and Urban Planning, 125: 222-233.
- Yao, H., Bowman, D. and Shi, W. (2006). Soil microbial community structure and diversity in a turfgrass chronosequence: Land-use change versus turfgrass management. Applied Soil Ecology, 34(2-4): 209-218.
- Zeng, N., Ding, Y. and Pan, J. (2008). Climate change-the Chinese challenge. Science, 319 (5864): 730-731.
- **Zhao, J.W., Shan, B.Q. and Yin, C.Q.** (2006). Effect of impervious surface on storm water runoff pollution in urban tourist attractions. Techniques and Equipment for Environmental Pollution Control, 7(12): 14-17.