

## SURVEY TO RECORD THE PEARL MILLET BLAST SEVERITY IN WESTERN RAJASTHAN

**Mukesh Kumar\*, Dama Ram<sup>1</sup>, J.R. Verma<sup>2</sup>, M.M. Sundria<sup>3</sup> and K.C. Bairwa<sup>4</sup>**

<sup>1</sup>*Department of Plant Pathology, CoA, Jodhpur (Agriculture University, Jodhpur, Rajasthan - 342304, India*

<sup>2</sup>*Department of Plant Pathology, ARS, Mandor (AU, Jodhpur), Rajasthan - 342304, India*

<sup>3</sup>*Department of Entomology, ARS, Mandor (AU, Jodhpur), Rajasthan - 342304, India*

<sup>4</sup>*Department of Agricultural Economics, CoA, Jodhpur (AU, Jodhpur), Rajasthan - 342304, India*

Email: [sihagmukesh6@gmail.com](mailto:sihagmukesh6@gmail.com)

*Received-24.09.2021, Revised-14.10.2021, Accepted-25.10.2021*

**Abstract:** A survey was conducted in major pearl millet growing districts of Western Rajasthan during *Kharif* 2020 to record the severity of blast disease. The highest pearl millet blast severity was recorded in Nagaur district (43.4%) Whereas, the minimum blast severity was recorded (16.6 %) in Pali district. The overall average disease severity of Western Rajasthan was (28.12%) based on total 125 fields surveyed in *kharif*2020.

**Keywords:** Survey, Pearl millet, Blast, Disease severity

### REFERENCES

- Anonymous** (2015). Annual Report; All India Co-ordinated Research Project on Pearl Millet 2014-15.
- DAC and FW** (2020). Fourth advance estimate, Department of Agriculture Cooperation & Farmers Welfare, Directorate of Economics and Statistics.
- Emchebe, A. M.** (1975). Some aspects of crop disease in Uganda UAFRO, Seren (unpublished).
- GoR** (2020). Commission rate of Agriculture, Jaipur, Rajasthan  
<http://www.agriculture.rajasthan.gov.in/content/dam/agriculture/Agriculture%20Department/agriculturalstatistics/CHTURTH%20%20AGRIM.pdf>. (accessed on 05.11.2020).
- Mehta, P. R., Singh, B. and Mathur, S. C.** (1953). A new leaf spot disease of bajra (*Pennisetum typhoides* Staph and Hubbard) caused by a species of *Pyricularia*. *Indian Phytopathol.*, 5: 140-143.
- Sharma, R., Upadhyaya, H. D., Manjunatha, S. V., Rai, K. N., Gupta, S. K. and Thakur, R. P.** (2013). Pathogenic variation in the pearl millet blast pathogen *Magnaporthe grisea* and identification of resistance to diverse pathotypes. *Plant Dis.*, 97(2): 189-195
- Singh, B. P. and Nainawatee, H. S.** (1999). Grain quality traits. In pearl millet breeding. Eds. Khairwal, I. S. Rai, K. N., Andrews, D. J. and Harinaryana, G. Oxford and IBH publishing Co. Pvt. Ltd., New Delhi, pp. 158-159.
- Thakur, R. P., Sharma, R. and Rao, V. P.** (2011). Screening Techniques for Pearl Millet Diseases. Information Bulletin No. 89. Patancheru 502324, Andhra Pradesh, India: International Crops Research Institute for the Semi-Arid Tropics. p.15.
- Wheeler, B.E.J.** (1969). An introduction of Plant diseases. John Willy & sons Ltd., London, p. 301.
- Wilson, J. P., Wells, H. D. and Burton, G. W.** (1989). Inheritance of resistance to *Pyricularia grisea* in pearl millet accessions from Burkino Faso and inbred Tift 85DB. *J. Heredity*, 80: 499-501
- Wilson, J. P., Wells, H. D. and Burton, G. W.** (1989). Inheritance of resistance to *Pyricularia grisea* in pearl millet accessions from Burkino Faso and inbred Tift 85DB. *J. Heredity*, 80: 499-501

\*Corresponding Author