

# EFFECT OF ZINC AND IRON APPLICATION ON YIELD AND ACQUISITION OF NUTRIENT ON MUSTARD CROP (*BRASSICA JUNCIA* L.)

Anuj Kumar, Satendra Kumar, Pradeep Kumar\* and Pramod Kumar

Department of Soil Science, College of Agriculture Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut 250 110

\*Email : [sehravat@gmail.com](mailto:sehravat@gmail.com)

**Abstract:** The field experiment was conducted on Pusa Bold variety of Mustard with 10 treatments in RBD in rabi season-2009-10 at Crop Research Centre of, Sardar Vallabhbhai Patel University of Agriculture and Technology; Meerut (U.P). Maximum primary branches (11.05), secondary branches (31.33), Silique per plant (545.35), number of seed per Silique (13.55), seed weight per plant 30.38 g and test weight (1000 seed weight, 6.50 g) were recorded, the biological yield was observed highest ( $114.80 \text{ q ha}^{-1}$ ) and the grain yield was also ( $23.40 \text{ q ha}^{-1}$ ) in T9{ 100 per cent NPK (RDF) + Zn @ 25 Kg  $\text{ha}^{-1}$  (B) + Fe @ 25 Kg  $\text{ha}^{-1}$  (B)}. The maximum Stover yield noticed  $91.40 \text{ q ha}^{-1}$  as compared to T1 (control) ( $40.82 \text{ q ha}^{-1}$ ), highest total nitrogen uptake by mustard crop, recorded  $97.87 \text{ kg/ha}$ , in case of phosphorus and potassium uptake by mustard crop was also observed  $21.82 \text{ kg/ha}$  and  $152.82 \text{ kg/ha}$ , respectively. The all over present investigation shows that the maximum yield attributes was found when zinc and iron was applied basal with recommended dose of fertilizers.

**Keywords:** Mustard, micronutrient, uptake  $\text{Kg ha}^{-1}$

## INTRODUCTION

India is one of the leading oil seed producing country in the world. Rapeseed and mustard are the main oil seed crops grown in rabi season in India. Oil seeds the second largest agricultural commodity after cereals in India. Its production was 7.20 tonnes from 6.3 m ha mainly confined with the states viz., Rajasthan, Uttar Pradesh, Haryana, Madhya Pradesh, Gujarat, West Bengal, Assam, Bihar, and Punjab. Among the states, Uttar Pradesh alone produces about 13.78 per cent of total mustard production from 14.03 per cent area, whereas, Rajasthan on top with 48.64 per cent production from 45.06 per cent area in India during 2008-09.

Iron is critical for chlorophyll formation and photosynthesis. Chlorophyll is the small "sun-panels" which the plants use to harvest energy from the sun and gives plants green pigment. Photosynthesis is the process during which the actual sun- rays are harvested. Iron is also used by enzymes to regulate transpiration in plants. This transpiration process allows nutrients to reach all parts of the plants. Without iron the above functions would not work.

Since these functions are essential for plant growth, iron is an essential element. So there is a need to focus on these nutrients, especially zinc as it is one of the most important micronutrient. While, applying with iron and NPK. Zinc deficient soil can be found throughout the world and are normally associated with low soil organic matter and alkaline of soil. Zinc deficiencies are corrected in most cases by applying a granular Zinc ( $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ ) fertilizer. Growth of winter crops in the soils is adversely affected due to reduction in zinc availability at low temperatures.

Although zinc can be applied as foliar application in emergency measure, greatest yields are obtained when it is applied to the soil. Soil application of zinc

is normally made at the seeding of crop. Sometimes, Zn deficiency appears due to the low amount of Zn as recommended and low temperature as required during the crop growth. Deficiency of zinc can also appear after seeding of the crop in soils with high phosphorus contents. Zinc sulphate improves phosphorus utilization and regulates plant growth and increase leaf size, promotes silking, hastens maturity and increase to test weight.

## MATERIAL AND METHOD

A field experiment was conducted during the rabi season of 2008-09 at the crop research center of SVBPUAT, Meerut. The soil was sandy loam with pH 8.36 and low in organic carbon 0.36per cent, available N (79.80), available K (165.30) and medium in available P (14.80)  $\text{kg ha}^{-1}$  and sufficient amount of Zn and Fe (0.49 and 12.25 ppm). The treatments T<sub>1</sub>-Control (without fertilizers), T<sub>2</sub>-100 per cent NPK @ 80,60,40  $\text{Kg ha}^{-1}$  (RDF), T<sub>3</sub>- 100 per cent NPK (RDF) + Zn @ 25  $\text{Kg ha}^{-1}$  (B), T<sub>4</sub>- 100 per cent NPK (RDF) + Zn (F), T<sub>5</sub>- 100 per cent NPK (RDF) + Fe @ 25  $\text{Kg ha}^{-1}$  (B), T<sub>6</sub>- 100 per cent NPK (RDF) + Fe (F), T<sub>7</sub>- 100 per cent NPK (RDF) + Zn @ 25  $\text{Kg/ha}$  (B) + Fe (F), T<sub>8</sub>- 100 per cent NPK (RDF) + Zn (F) + Fe @ 25  $\text{Kg ha}^{-1}$  (B), T<sub>9</sub>- 100 per cent NPK (RDF) + Zn @ 25  $\text{Kg ha}^{-1}$  (B) + Fe @ 25  $\text{Kg ha}^{-1}$  (B) and T<sub>10</sub> - 100 per cent NPK (RDF) + Zn (F) + Fe (F) were laid out in RBD in the three replications. The calculated quantity of Zn and Fe was applied as basal as well as foliar at first and second irrigation as per treatment 30 and 60 DAS, respectively. Indian mustard variety Pusa Bold was sown at 30 cm row spacing using 4 kg seed  $\text{ha}^{-1}$ .

## RESULT AND DISCUSSION

### Effect on growth and yield and yield attribute of mustard

The data on growth parameters and yield and yield attribute of mustard are presented in Table 1 and 2. Various plant growth parameter of mustard crop are affected by varying the method of application of micronutrients during the crop season. The plant height was significantly influence by the different method of zinc and iron application at all the growth stages. Plant height is an index of good vegetative growth. The maximum plant height 98.25 cm at 60 DAS and 196.25 cm at harvest was observed in T<sub>9</sub> [100 per cent NPK (RDF) + Zn @ 25 Kg ha<sup>-1</sup> (B) + Fe @ 25 Kg/ha (B)] treatment, i.e. 36.45 and 26.20 per cent increase in plant height were observed at 60 and 147 DAS, respectively (Table 2). Primary branches increase with advancement of crop age the maximum branches was observed in T<sub>9</sub> (11.05) at 60 DAS significantly, when zinc and iron applied basal and foliar (Table 1). The similar results were also reported by Meena *et al.* (2006), Chaudhary *et al.* (2007), Yadav *et al.* (2007) and Ravi *et al.* (2008). Maximum number of secondary branches at 60 DAS (31.33) and at harvest 147 DAS (34.67) were recorded in T<sub>9</sub> [100 per cent NPK (RDF) + Zn @ 25 Kg/ha (B) + Fe @ 25 Kg/ha (B)]. (Table 1) significantly increase in number of branches per plant as compared to T<sub>1</sub> attributed to increase in absorption and translocation of assimilation and stimulation graphical and lateral meristems to grow result supported by Husain and Kumar (2006).

The numbers of Siliqua per plant at harvesting highest (545.35), number of seed per Siliqua was recorded maximum in T<sub>9</sub> (13.55). The highest length of Siliqua found 5.20 cm (Table- 2), treatment in which recommended dose of fertilizer were applied along with zinc and iron application as basal. The significant increase in number of Siliqua per plant and number of seed per Siliqua as compared to T<sub>1</sub>(Control). It is evident that zinc and iron element play an important role in plant, similar results were reported by Sudhakar *et al.* (2002) and Husain and Kumar (2006). Significantly result was observed, when zinc and iron was applied basal along with RDF (T<sub>9</sub>) in seed weight per plant and 1000 seed weight (6.50g) the result was supported by Sudhakar *et al.* (2002), and Zizala *et al.* (2008).

### Yield attributes

The grain yields of mustard in different treatments are significant; when zinc and iron are applied alone and with combination significantly increased the grain yield. The highest grain yield was recorded in

T<sub>9</sub> (23.40 q/ha) followed by T<sub>7</sub> and T<sub>8</sub> gave grain yield 22.14 and 20.45 qha<sup>-1</sup>. i.e. 145, 132 and 115 per cent increased in grain yield by T<sub>9</sub>, T<sub>7</sub> and T<sub>8</sub>, respectively over control. (Table 2). Similar observation were also recorded by Saxena *et al.* (2005), Kumar *et al.* (2006), Meena *et al.* (2006), Chaudhary *et al.* (2007), Jat and Mehra (2007), Yadav *et al.* (2007) and Ravi *et al.* (2008). Similarly the Stover yield was recorded significantly maximum in T<sub>9</sub> (91.40 q ha<sup>-1</sup>) followed by T<sub>7</sub>, T<sub>8</sub> and T<sub>10</sub> gave values 87.74, 79.25 and 75.82 q ha<sup>-1</sup>, respectively. The stover yield was observed 123, 114, 94 and 86 per cent higher as compared to T<sub>1</sub> (control). These results are supported by the findings of Saxena *et al.* (2005), Chaudhary *et al.* (2007), Jat and Mehra (2007) and Chandra and Khandelwal (2009).

### Nutrient uptake

The data on N, P and K uptake by grain of mustard are presented in Table 3. Total uptake of NPK by mustard crop was maximum recorded with RDF along with zinc and iron applied as basal. The total nitrogen uptake was recorded significantly highest in T<sub>9</sub>. Total phosphorus uptake was found maximum in T<sub>9</sub> followed by T<sub>7</sub> over control, and the total potassium uptake by mustard crop was also in highest in T<sub>9</sub> over the maximum per cent in increase of nutrient was recorded 191, 236 and 222 per cent NPK, respectively over control. The result are supported by Malewar *et al.* (2001), Giri *et al.* (2003), Kumar *et al.* (2006), Meena *et al.* (2006), Chaudhary *et al.* (2007), Jat and Mehra (2007), Ravi *et al.* (2008), Zizala *et al.* (2008) and Chandra and Khandelwal (2009).

## CONCLUSION

It is concluded from that investigation the application of recommended dose of fertilizer (100per cent NPK) @ 80: 40: 40 recorded better grain yield (11.90 q ha<sup>-1</sup>) of mustard crop. The highest grain yield (23.40 q ha<sup>-1</sup>) was obtained in the treatment consisting the basal application of zinc and iron along with 100per cent nitrogen, phosphorus and potash (Recommended dose of fertilizer) (T<sub>9</sub>). The addition of Zn and Fe as basal @ 25 Kg/ha along with 100per cent NPK (RDF) prone super ion to foliar application of Zn and Fe along with 100per cent NPK (RDF) in terms of yield and other parameters of mustard crop. On an average highest total uptake of NPK recorded 97.54, 21.82, and 152.82 Kg ha<sup>-1</sup>, respectively in treatment T<sub>9</sub> followed by T<sub>7</sub>. Similarly maximum total uptake of zinc and iron were recorded 251.76 and 2314.53 g ha<sup>-1</sup> respectively in T<sub>9</sub> followed by T<sub>7</sub>.

**Table 1:** Effect of zinc and iron application on number of primary and secondary branches/plant in mustard.

Treatment	primary branches/plant		secondary branches/plant	
	30 DAS	60 DAS	60 DAS	At harvest
T <sub>1</sub>	1.50	4.75	11.33	13.33
T <sub>2</sub>	1.58	5.75	23.00	25.33
T <sub>3</sub>	1.61	7.97	23.33	25.67
T <sub>4</sub>	1.56	6.65	20.00	23.33
T <sub>5</sub>	1.46	7.35	22.33	25.00
T <sub>6</sub>	1.50	6.15	20.00	22.33
T <sub>7</sub>	1.57	10.25	24.67	27.67
T <sub>8</sub>	1.74	8.50	22.00	24.33
T <sub>9</sub>	1.76	11.05	31.33	34.67
T <sub>10</sub>	1.72	8.17	23.67	27.00
S Em ( $\pm$ )	0.10	0.30	1.68	1.61
CD (P=0.05)	N.S.	0.91	5.04	4.81

**Table 2:** Effect of zinc and iron application on yield and yield attributes of mustard crop.

Treatment	Grain yield (q ha <sup>-1</sup> )	Stover yield (q ha <sup>-1</sup> )	Biological yield (q ha <sup>-1</sup> )	Plant height (cm)	No. of silique/ plant	No. seed/ silique	Length of silique (cm)	Seed weight/ plant (g)	1000- seed weight (g)
T <sub>1</sub>	9.53	40.82	50.35	155.50	295.25	7.95	3.50	9.45	5.20
T <sub>2</sub>	11.90	56.52	68.42	160.67	390.60	8.40	4.02	12.10	5.40
T <sub>3</sub>	17.94	74.14	92.08	177.25	485.40	10.15	4.62	17.10	5.65
T <sub>4</sub>	13.90	71.43	85.33	169.25	460.45	9.45	4.40	14.11	5.55
T <sub>5</sub>	14.00	74.14	88.14	170.25	478.75	9.65	4.55	15.20	5.80
T <sub>6</sub>	12.20	66.91	79.11	165.00	415.15	8.90	4.36	13.50	5.85
T <sub>7</sub>	22.14	87.74	109.88	188.25	522.50	12.20	4.90	28.78	6.35
T <sub>8</sub>	20.45	79.25	99.70	182.00	505.75	11.95	4.85	25.32	6.20
T <sub>9</sub>	23.40	91.40	114.80	196.25	545.35	13.55	5.20	30.38	6.50
T <sub>10</sub>	19.28	75.82	95.10	180.25	490.65	10.75	4.75	20.51	6.00
S Em $\pm$	0.40	0.51	0.59	2.34	4.00	0.16	0.16	0.29	0.14
CD (P=0.05)	1.19	1.52	1.77	7.01	11.97	0.48	0.48	0.87	0.40

**Table 3:** Effect of zinc and iron application on Total NPK uptake (kg/ha) by mustard crop.

Treatments	Nitrogen uptake (kg/ha)	Phosphorus uptake(kg/ha)	Potassium uptake (kg/ha)
T <sub>1</sub>	23.53	6.47	47.25
T <sub>2</sub>	34.71	11.21	90.21
T <sub>3</sub>	47.69	17.10	92.95
T <sub>4</sub>	44.03	18.65	116.00
T <sub>5</sub>	43.39	15.62	119.01
T <sub>6</sub>	38.86	13.58	105.94
T <sub>7</sub>	60.46	19.89	148.83
T <sub>8</sub>	55.93	17.88	134.34
T <sub>9</sub>	67.19	21.82	152.82
T <sub>10</sub>	54.28	17.69	124.09
S Em $\pm$	6.43	0.26	10.55
CD (P=0.05)	19.25	0.78	31.59

## REFERENCES

- Chandra, D. and khandelwal, R.B.** (2009). Effect of zinc and phosphorus on yield, nutrient uptake and oil content of mustard grown on the gypsum –treated sodic soil. *J. Indian soci. Soil Sci.*, 57 (1): 66-70.
- Chaudhary, H.C., Khan, S. and Uttam, S.K.** (2007). Effect of N, P, S and Zn nutrition and moisture conservation practices on nutrient uptake, quality and yield of rainfed Indian mustard in eroded soil. *Indian Agriculture*, 53 (3/4): 125-130.
- Giri, M.D., Hamid, A., Giri, D.G., kumar,R.P. and Sajid, M.** (2003). Effect of irrigation and sources of iron on quality and uptake of nutrients of mustard. *J. soils and crops*, 13 (1):131-134.
- Husain, M.F.and Kumar,R.** (2006). Influence of sowing dates application of zinc on the performance of mustard in South-West semi arid zone of Uttar Pradesh. *International J. Agriculture Sci.*, 2(2):601-604.
- Jat, J.R. and mehra,R.K.** (2007). Effect of sulphur and zinc on yield, macronutrient content in and uptake by mustard Haplustepts. *J. Indian soci.soil sci.*, 55 (2): 190-195.
- Kumar,N.,Singh,S.and Singh,V.** (2006). Effect of Iron and Sulphur levels on yield, oil content. and their uptake by on Indian mustard (*Brassica juncea*). *Indian J.Agronomy*, 51(1):63-64.
- Malewar,G.V., Kate, S.D., Waikar, S.L. and Isnail, S.** (2001). Interaction effects of zinc and boron on yield, nutrient uptake and quality of mustard (*Brassica juncea* L.) on a typic Haplustert. *J. Indian soci. Soil sci.*, 49(9):763-765.
- Meena, M. C., Patel, K. P. and Rathod, D. D.** (2006). Effect of Zinc, Iron and Sulphur on mustard in loamy sand soil, *Indian J. Fertilizer*, 2(5):55-58.
- Ravi, S., channel, H.T.,Hebsur, N.S.,Patil,B.N.and Dharmatti, P.R.** (2008). Effect of sulphur, zinc and iron nutrition on growth, yield, nutrient uptake and quality of safflower (*carthamus tinctorius* L.). *Karnataka J.agric.sci.*, 21 (3): 382-385.
- Saxena,K.K.,Kumar,A.and Verma,H.R.** (2005). Growth,yield and oil content of mustard (*Brassica juncea* L.) as influenced by application of phosphorus and iron. *Farm Sci J.* 14(1): 97-98.
- Sudhakar,P.C.,Chandel,R.S. and Singh,K.** (2002). Effect of sulphur, iron and silicon on the growth and yield of irrigated mustard. *Annals of Agricultural Res.*, 23 (3): 483-485.
- Yadav,R.b.,Singh,R.V., Singh, H.R. and yadav, H.S.** (2007). Effect of different levels of zinc and sulphur on Indian mustard (*Brassica juncea* L.). *crop res. Hisar*, 33 (43): 74-76.
- Zizala,V.J.,jadav,N.B. and Gorfad,P.S.** (2008). Effect of sulphur and zinc on yield, quality and its concentration on mustard. *Asian J. soil sci.*, 3 (1): 173-177.