

# FLORA OF JHAJJAR DISTRICT, HARYANA

**Jai Bhagwan Singh**

*Department of Botany, Meerut College, Meerut*

**Abstract:** The present work comprises of 731 species belonging to 459 genera and 114 families of Angiosperms. Of these 611 species under 371 genera and 93 families belong to Dicotyledones and 120 species under 88 genera and 21 families belong to monocotyledones. The ratio of families belonging to monocots and dicots were 1:4.46; of genera 1:4.21 and of species 1:5.09. The ratio of genus to species is 1:1.64.

**Key words:** Family, Flora, Genus, Jhajjar district.

## INTRODUCTION

Diverse climate and habitat of India provide favourable conditions for colonisation of a variety of plant species. The flora of India thus show great diversity. It is one of the 12 mega diversity centre of the world with over 17,000 species of flowering plants. There are a large number of endemic species in the Himalaya and Peninsular India.

District Jhajjar came into being as a result of bifurcations of Rohtak district on 15th July 1997. Inhabited about eight hundreds years ago, Jhajjar city has many pages of history engulfed in it. At the time of Gori's attack in 1191 AD the area was a deserted forest. There was a town named Malokan in the east of present Jhajjar city mostly inhabited by Jats. Jhajjar district was named to commemorate the brave Jat Chajju of Malokan village who fought bravely to defend the country against Gori's.

The district is bounded by Bhiwani in the west, Delhi in the east, Gurgoan and Rewari in the South and Rohtak and Sonapat districts in the north respectively. The district lies at 28° 37'N latitude and 76° 39' longitude. The district is divided into three tehsils - Jhajjar, Bahadurgarh and Beri and comprises of five blocks - Jhajjar, Bahadurgarh, Sahlawas, Matanhail and Beri. The district is well known for its historical places like, Jhan Ara Bagh, Gurukul Jhajjar Museum and Bhimeshwar Devi Temple.

John Forbes Royle (1799-1858), an English armyman, was the pioneer plant explorer of Northern India and Himalayan Mountains. He succeeded Govan as the Curator of the new Botanical Gardens at Saharanpur in 1823. Stewart (1869) studied the plants of Punjab and published under the title Punjab Plants. Flora of Upper Gangetic Plain and of the Adjacent Siwalik Hills and Sub-Himalayan tracts by Duthie (1903-1929). Flora of British India by Hooker (1872-1879). Flora of Delhi by J.K. Maheshwari (1963, 1966) are some other important publications on the flora of North-West India. A Catalogue of the Plants of the Punjab and Sindh (Atchison, 1869); Forest flora of North-West and Central India (Brandis,

1874); Grasses of North. West India - Indigenous and Cultivated (Duthie, 1883); Field and Garden Crops of North-West Provinces and Oudh. (Duthie & Fuller, 1882-1893); Forest Flora for the Punjab with Hazara and Delhi (Parker, 1918); the Flora of India Desert (Jodhpur and Jaisalmer) (Blatter *et al.*, 1918-21); Forest Flora of Jaipur State (George, 1937); Common Grasses of the United Provinces (Bor, 1947); A contribution to the Flora of Punjab Plains and associated hill regions (Sabnis, 1940-41); Grasses of Jaipur (Gandhi *et al.*, 1961); Some new records for the Punjab Plain (Nair, 1963 a.b., 1964, 1966); Flora of Rajasthan, West of Aravalis (Puri *et al.*, 1964), Flora Punjab Plains (Nair, 1978); Flora of Indian Desert (Bhandari, 1978); Flora of North-East Rajasthan (Sharma *et al.*, 1979) and Flora of North-East Haryana (Jain *et al.*, 1982).

Flora of Jhajjar in Haryana has not been studied and to fill up this gap a comprehensive study of the Flora of Jhajjar was undertaken.

## MATERIALS AND METHODS

The present work is based on intensive study of plants in the Jhajjar district for four consecutive years. The collection of plants was followed by their further study in the laboratory and their identification were finally confirmed at the herbaria of Northern Circle of Botanical Survey of India and Forest Research Institute, Dehradun.

### A. Collection

All efforts were made to make excursions periodically in different parts of the district at regular intervals. As an outcome of this, it was possible to collect most of the plants in vegetative, flowering and fruiting stages from different localities and habitats. Intensive exploration was done in the areas with little biotic interference. Particular attention was paid to the areas adjoining rivers, canals and swamps.

In the course of the day's excursion, field observations including habit and habitats, soil type, flower colour, time

of flowering and fruiting, relative abundance were noted in a field note book. Complete plants or good size twigs with flowering and fruiting stages were gathered. All the plants were tagged with a number.

### B. Preservation

The plants were pressed in old newspapers and blotting sheets for dehydration in strong plyboard presses in the field and later transferred to heavy metal plant press for perfect pressing. Corrugated sheets were used in between the newspapers sheets which permitted circulation of air. The specimens were changed to fresh sheets after an interval of 24 hours to 2-3 days depending on the weather conditions until the specimens were completely dry. The number and frequency of change was altered according to dryness of climate and fleshiness of tissues, for succulents the interval of change was frequent and for grasses it was short. The succulent plants were boiled in water before pressing. Dissected parts and loose parts such as large fruits and seeds were placed in paper packets pasted to the mounting sheet. Before mounting, the dehydrated specimens were poisoned by dipping them in saturated solution of mercuric chloride in ethyl alcohol. Nephthalene balls and periodical fumigation with formaldehyde were used as insect repellent during storage of herbarium sheets in almirahs. The dried and poisoned specimens were mounted on white herbarium sheets of standard size (43×28 cm). Fevicol (synthetic resin adhesive) was used as pasting material for plants which sticks fast and also minimizes the chances of insects attack.

### C. Identification

The identification of plants was done with the help of Flora of British India (Hooker, 1872-1897), Flora of Upper Gangetic Plain and of the Adjacent Siwalik and Sub-Himalayan Tracts (Duthie, 1903-1929), Supplement to the Duthies Flora of Upper Gangetic Plain (Raizada, 1976), Herbaceous Flora of Dehradun (Babu, 1977), Flora of Delhi (Maheshwari, 1963) and other floras and monographs. For grasses the following floras were used; Grasses of Upper Gangetic Plain (Raizada and Jain, 1957, 1964); Grasses of Burma, Ceylon, India and Pakistan (Bor, 1960) and A list of Grasses of North-Western India (Duthie 1883). For cultivated plants, Manual of Cultivated Plants (Bailey, 1949) and J.F. Duthi's (1883) Indigenous and Cultivated Plants of Roorkee were consulted.

### D. General Plan Of Present Work

In the present work for the sake of convenience, Benthem and Hooker's system of classification (1862-1883) was followed. Splitting of families was done following, Hutchinson (1959) and Airy Shaw's (1973) Genera in each family and species in each genus were arranged in alphabetical order for the convenience of reference.

Artificial keys are provided for genera within a family and species within a genus. The most convenient key is dichotomous that is based on successive choices between two contrasting alternative characters. The efforts are made to make artificial key as simple as possible without ignoring prime characters.

An effort has been made to make the plant nomenclature upto date as far as possible after consulting available literature. The valid name of a species is followed by a reference to "Flora of British India", "Flora of Upper Gangetic Plain" and other recent literature. Basionyms and synonyms for species are also provided.

A short description is provided for each species. The description is followed by habit, abundance, flowering period, Local names and local uses. In general, it has been noticed that only such plants which are useful to local people have Local names.

## DESCRIPTION

### Statistical Analysis of the Flora

During the floristic studies of Jhajjar district the total number of species recorded was 731 belonging to 459 genera and 114 families of Angiosperms. Of these 601 species under 371 genera and 94 families belong to dicotyledones and 120 species representing 88 genera and 94 families belong to dicotyledones and 120 species representing 88 genera and 21 families belong to Monocotyledones. The ratio of families belonging to monocotyledones and dicotyledones was 1:4.46, of genera 1:4.21 and of species 1:5.09. The ratio of genera to species was 1:1.64 as compared to 1:1.63 for Delhi State, 1:2.2 for Upper Gangetic Plain and 1:7 for British India and 1:6 for India alone (Chatterjee, 1939). The table-1 shows the statistical data for various categories of flora.

**Table 1. Statistical analysis**

	Dicots		Monocots		Total
	Number	%age	Number	%age	
<b>Families</b>	93	81.73	21	18.26	114
<b>Genera</b>	371	80.82	88	19.17	459
<b>Species</b>	611	83.53	120	16.41	731

The analysis of flora indicates that the ten dominant families comprise nearly half the total plants recorded from the district. These families with their number of species, and genera in order of dominance are given below in Table-2.

**Table 2. Ten dominant plant families of Jhajjar district**

S. No.	Name of Family	Number of genera	Number of species
1.	Fabaceae	45	75
2.	Poaceae	35	57
3.	Asteraceae	35	44
4.	Malvaceae	12	24
5.	Acanthaceae	12	21
6.	Convolvulaceae	06	20
7.	Scrophulariaceae	12	19
8.	Solanaceae	08	19
9.	Verbenaceae	11	17
10.	Boraginaceae	11	17

The analysis shows that the monocotyledones were less dominant and poorly represented except for the families Poaceae and Cyperaceae. Out of 120 species of monocotyledones 72 species belong to two families poaceae (57 species) and cyperaceae (15 species) while the remaining 659 species belong to 113 families.

In order to find out the relation of the present floristic investigation with other neighbouring and important floras a comparative list of 10 dominant families based on the number of species is given below (Table-3).

**Table 3. Ten dominant families of district Jhajjar, Delhi State, Upper Gangetic Plain and British India**

Sl. No.	District Jhajjar	Delhi State (After Maheshwari, 1963)	Upper Gangetic Plain (After Duthie, 1903-1922)	British India (After Hooker, 1904)
1.	Fabaceae	Poaceae	Poaceae	Orchidaceae
2.	Poaceae	Fabaceae	Fabaceae	Fabaceae
3.	Asteraceae	Asteraceae	Cyperaceae	Poaceae
4.	Malvaceae	Cyperaceae	Asteraceae	Rubiaceae
5.	Acanthaceae	Acanthaceae	Scrophulariaceae	Euphorbiaceae
6.	Scrophulariaceae	Euphorbiaceae	Malvaceae	Acanthaceae
7.	Verbenaceae	Convolvulaceae	Acanthaceae	Asteraceae
8.	Boraginaceae	Malvaceae	Euphorbiaceae	Cyperaceae
9.	Euphorbiaceae	Amaranthaceae	Convolvulaceae	Lamiaceae
10.	Convolvulaceae	Scrophulariaceae	Lamiaceae	Urticaceae

The above comparison of 10 dominant families shows that the families Poaceae and Fabaceae occupy first two positions in the floras of district Jhajjar, Delhi State and Upper Gangetic plains whereas Asteraceae, Acanthaceae and Euphorbiaceae are amongst the ten dominant families in all the four floras. This shows some similarity in the pattern of the floras of district Jhajjar, Delhi State, Upper Gangetic Plain and British India.

### ACKNOWLEDGEMENTS

I feel great pleasure in expressing my deep sense of gratitude to my esteemed teacher Late Dr. H. Singh, Reader, Department of Botany, Meerut College, Meerut for his painstaking guidance, generous help, stimulating comments and suggestions which enabled me to bring this work to its present form.

### REFERENCES

**Airy Shaw, H.K.** (1952). Note on Taxonomic position of *Nyctanthes* Linn. and *Dimetra* Kerr. *Kew Bull.* 271-272.

**Airy Shaw, H.K.** (1973). *Willi's Dictionary of Flowering Plant and Ferns*. Ed. B. London.

**Aitchison, J.E.T.** (1869). *A catalogue of the Plants of the Punjab and Sindh*, London.

**Andrews, F.W.** (1950-56). *The flowering Plants of the Anglo-Egyptian Sudan*. Scotland. 3 Vols.

**Ansari, M.Y.** (1984). *Asclepiadaceae : Genus Ceropogia* in Fascicles of Flora of India, No. 16, *Botanical Survey of India, Howrah*.

**Babu, C.R.** (1977). *Herbaceous Flora of Dehradun*. CSIR, New Delhi.

**Bamber, C.J.** (1966). *Plants of the Punjab*. Lahore.

**Bailey, L.H.** (1949). *Manual of Cultivated Plants*. (Ed.2), New York.

**Bentham, G. and Hooker, J.D.** (1862-1883). *Genera Plantarum*. Vol. 1-3, London.

- Benthall, A.P.** (1946). The Trees of Calcutta and its neighbourhood. Calcutta.
- Bhandari, M.M.** (1978). Flora of Indian Desert. *Scientific Publisher, Jodhpur*.
- Bhattacharya, U.C.** (1963a). A contribution to the Flora of Mirzapur-1. Some new records for the district and for Upper Gangetic Plain. *Bull. Bot. Sur. India*, **5**:59-62.
- Bhattacharya, U.C.** (1964a). A contribution to the Flora of Mirzapur-II, *Bull. Bot. Sur. India*, **6**:191-210.
- Bhattacharya, U.C. and Malhotra, C.L.** (1964b). Botanical exploration in Hamirpur district, U.P. *Bull. Bot. Sur. India*, **6**(1):25-41.
- Blatter, E. and Hallberg, F.** (1917-1918). A revision of the Indian species of *Rotala* and *Ammannia*. *J. Bom. Nat. Hist. Soc.* **25**:701-722; **26**:210-217.
- Blatter, E. and Hallberg, F.** (1918-1921). The Flora of Indian Desert (Jodhpur and Jaisalmer). *Bomb. Nat. Hist. Soc.* **26**:218-240, 525-551, 811-818, 968-987; *ibid* **27**:40-47, 270-279, 506-519.
- Blatter, E. and McCann.** (1930-1935). Revision of the Flora of Bombay Presidency. *J. Bom. Nat. Hist. Soc.* **34**:291-306, 623-637, 877-900; *ibid* **36**:524-537, 781-785; *ibid* **37**:15-35, 255-277, 532-548, 764-779.
- Bor, N.L.** (1947). Common Grasses of the United Provinces. *Ind. For. Rec. (Botany)*, **2**:1-220.
- Bor, N.L.** (1952). The Genus *Poa* Linn. in India. *J. Bom. Nat. Hist., Soc.*, **50**:787-838.
- Bor, N.L.** (1960). The Grasses of Burma, Ceylon, India and Pakistan. *Pregamon Press, London*.
- Brandis, D.** (1966). Indian Trees (Repr. Ed. 1971). Bishen Singh and Mahendra Pal Singh, Dehradun.
- Brenan, J.P.M.** (1957). Notes on Mimosidaceae III. *Kew Bull.*, **1**:75-96.
- Chakravarty, H.L.** (1982). Cucurbitaceae: in Fascicles of Flora of India, No. 11, *Botanical Survey of India, Howrah*.
- Chatterjee, D. and Bharadwaja, R.C.** (1956). Revision of the Scrophulariaceae of Upper Gangetic Plain and Adjoining areas. *Bull. Bot. Soc. Bengal*, **9**:127-153.
- Chopra, R.N.; Nayar, S.L. and Chopra, I.C.** (1956). Glossary of Indian Medicinal Plants. CSIR, New Delhi.
- Collett, H.** (1921). *Flora Similensis* (Reprinted, 1971). Bishen Singh and Mahendra Pal Singh, Dehradun.
- Croizat, Leon** (1940). Notes on Indian Euphorbiaceae *Croton bonplandianum* (*C. sparsiflorum*) and *Euphorbia perbracteata*. *J. Bom. Nat. Hist. Soc.*, **41**:573-576.
- Debnath, H.S. and Nayar, M.P.** (1984). Papaveraceae and Hypecoaceae : In Fascicles of Flora of India. No. 17, *Botanical Survey of India, Howrah*.
- Deshpande, U.R.** (1984). Poaceae : Tribe-Andropogoneae: In Fascicles of Flora of India.
- Dixit, S.N.; Verma, S.D. and Srivastava, T.N.** (1966). Additions to the rainy season weeds of Gorakhpur. *Proc. Nat. Acad. Sci., India*, **36B**:149-156.
- Duthie, J.F.** (1883). A list of Grasses of North-Western India. Indigenous and cultivated. Roorkee.
- Duthie, J.F. and Fuller, J.B.** (1882-1893). Field and Garden Crops of the North-West Provinces and Oudh. Roorkee.
- Duthie, J.F.** (1885). Report on an examination of the indigenous grasses and other fodder yielding plants growing on the Hissar Bir Land. Roorkee.
- Duthie, J.F.** (1888). The Fodder Grasses of Northern India. Roorkee.
- Duthie, J.F.** (1903-1929). Flora of the Upper Gangetic Plain and of the Adjacent Siwalik and sub-Himalayal tracts (Ed.1), 3 Vol., Calcutta.
- Engler, A and Prantl., K.** (1887-1915). Die Naturlichen Pflanzefamilier, etc.
- Fyson, P.F.** (1922). The Indian species of *Eriocaulon*. *J. Indian Bot. Soc.*, **3**:12-18.
- Gandhi, S.M.; Bhargawa, P.D. and Bhatnagar, M.P.** (1961). Grasses of Jaipur. *Proc. Nat. Acad. Sci. India B* **31**-2:183-192
- Ghora, C. and Panigrahi, G.** (1984). Rosaceae: Genus *Prunus* : in Fascicles of Flora of India. No. 18, *Botanical Survey of India, Howrah*.

- Gupta, B.L.** (1928). Forest Flora of Chakrata, Dehradun and Saharanpur Forest Division. Bishen Singh and Mahendra Pal Singh, Dehradun.
- Gupta, R.K.** (1969). Common Grasses of Gorakhpur. *Ind. For.*, **95**:324-329.
- Haines, H.H.** (1921-1924). The Botany of Bihar and Orissa, London. 6 parts.
- Hajara, P.K.** (1983). Linaceae and Ixonanthaceae : in Fascicles of Flora of India.
- Hooker, J.D.** (1872-1897). *The Flora of British India*. Vol. 1-7, London.
- Husain, W.** (1967). Some new records of plants for the Upper Gangetic Plain. *Ind. For.*, **93**:582-585.
- Hutchinson, J.** (1959). The Families of Flowering Plants. Ed. 2, Vol. 1-2, Oxford.
- Jain, R.K.** (1958). Hitherto unrecorded occurrence of *Datura innoxia* Mill. in the Upper Gangetic Plain and Rajasthan desert. *Curr. Sci.*, **27**:395.
- Jain, S.K. and Rao, R.R.** (1977). *A Handbook of Field and Herbarium Methods*. Today and Tomorrow's Printers and Publishers, New Delhi.
- Jain, S.P.; Singh, J.S. and Verma, D.M.** (1982). Flora of North-East Haryana (India). *J. Econ. Tax. Bot.*, **3**:152-176.
- Kanjilal, P.C.** (1933). Forest Flora of Pilibhit, Oudh, Gorakhpur and Bundelkhand, Allahabad.
- Kanjilal, P.C.** (1966). A Forest Flora for the Plains of Uttar Pradesh. Part II & III, Lucknow.
- Kundu, B.C. and Anima De** (1968). Taxonomic position of the genus *Nyctanthes*. *Bull. Bot. Sur. India*, **10**:397-408.
- Lawrence, G.H.M.** (1951). Taxonomy of vascular plants. New York.
- Maheshwari, J.K.** (1957). Some name changes in the Flora of India. *J. Bomb. Nat. Hist. Soc.* **54**:884-805.
- Maheshwari, J.K.** (1961). On the Identity and Nomenclature of some Indian Plants. *Bull. Bot. Surv. Ind.* **3**:91-94.
- Maheshwari, J.K.** (1963). The Flora of Delhi. CSIR, New Delhi.
- Maheshwari, J.K.** (1964). *Alternanthera philoxeroides* (Mart.) Griseb. A new record for India. *Bull. Bot. Sur. India*, **6**: 313-314.
- Maheshwari, J.K.** (1966). Illustrations to the Flora of Delhi. New Delhi.
- Malhotra, J.K.** (1966). New distributional record of plants from Upper Gangetic Plain. *Bull. Bot. Sur. India*, **8**: 7-78.
- Melville, R.** (1958). Notes on *Alternanthera*. *Kew Bull.*, **1**: 171-175.
- Mukerjee, S.K.** (1940). A revision of the Labiatae of the Indian Empire. *Rec. Bot. Sur. India*, **24**(1): 1-228.
- Mukerjee, S.K.** (1945). A revision of the Indo-Burmese species of *Lindernia allioni*. *J. Ind. Bot. Soc.*, **24**: 127-134.
- Mukerjee, S.K.** (1953). Vegetation of Delhi Ridge. *J. Bom. Nat. Hist. Soc.*, **51**: 438-465.
- Murty, Y.S. and Singh, V.** (1961). New plant records for the Upper Gangetic Plain from Meerut and its neighbourhood. *Proc. Nat. Inst. Sci. India*, **27B** (1): 13:17.
- Nair, N.C.** (1978). Flora of Punjab Plains. *Rec. Bot. Sur. India*. **21**(1): 1-326.
- Nair, N.C. and Nair, V.J.** (1963). Some plant records for the Punjab Plain. *Bull. Bot. Sur. India*, **5**(3-4): 219-222.
- Nair, N.C.** (1964a). Some new records for the Punjab plain II, *Bull. Bot. Sur. India*, **6**(1): 69-71.
- Nair, N.C. and Nair, V.J.** (1966). New plant records for the Punjab Plain IV. *Bull. Bot. Sur. India*, **8**(3-4): 351-353.
- Nair, N.C. and Nair, V.J.** (1977). Present status and future strategies of Floristic studies in India. The Upper Gangetic Plain. *Bull. Bot. Sur. India*, **19**(1-4):25-32.
- Pal, B.P.** (1972). *Beautiful Climbers of India*. ICAR, New Delhi.
- Parker, R.N.** (1918). A Forest Flora of the Punjab with Hazara and Delhi. Lahore, 3rd Edn., 1956.
- Parker, R.N. and Turrill, W.B.** (1929). Flora of the Upper Gangetic Plain and the Adjacent Siwalik and Sub-Himalayan Tracts—Palmae to Cyperaceae. Central Publication Branch, Calcutta.

- Puri, G.S.; Jain, S.K.; Mukherjee, S.K.; Sarup, S. and (Miss) Kotwal, N.N.** (1964). Flora of Rajasthan, West of Aravalis. *Rec. Bot. Surv. Ind.* **19**(×1):1-59.
- Raizada, M.B.** (1931). Contribution to Duthie's Flora of Upper Gangetic Plain from the neighbourhood of Dehradun. *J. Indian Bot. Soc.*, **10**: 155-158.
- Raizada, M.B.** (1935). Recently introduced or otherwise imperfectly known plants from the Upper Gangetic Plain. *J. Indian Bot. Soc.*, **14**: 339-348.
- Raizada, M.B.** (1936). Recently introduced or otherwise imperfectly know plants from the Upper Gangetic Plain. *J. Indian Bot. Soc.*, **15**: 149-167.
- Raizada, M.B.** (1939). Recently introduced or otherwise imperfectly known plants from the Upper Gangetic Plain. *J. Indian Bot. Rec. (NS) Bot.*, **1**: 223-235.
- Raizada, M.B.** (1950). New or noteworthy plants from the Upper Gangetic Plain. *Ind. For.*, **76**: 489-497.
- Raizada, M.B.** (1954). Grasses of the Upper Gangetic Plain and some aspects of their ecology. *Ind. For.*, **80**: 24-26.
- Raizada, M.B.** (1958). Name changes in common Indian plants. *Ind. For.*, **84**:467-538.
- Raizada, M.B.** (1958). New plants records for the Upper Gangetic Plain *Proc. Nat. Inst. Sci. India*, **24B** : 198-208.
- Raizada, M.B.** (1959). Name changes in common Indian Grasses. *Ind. For.*, **85**: 473-509.
- Raizada, M.B.** (1976). Supplement to Duthie's Flora of the Upper Gangetic Plain and of the Adjacent Siwalik and Sub-Himalayan Tracts. Bisen Singh and Mahendra Pal Singh, Dehradun.
- Raizada, M.B. and Jain, S.K.** (1964). Grasses of the Upper Gangetic Plain. Part II (Poaicoideae). *Indian For. Rec. (NS) Bot.*, **5**(7): 151-226.
- Raizada, M.B. and Jain, S.K.** (1966). Grasses of the Upper Gangetic Plain-Poaicoideae. *Indian For.*, 637-642.
- Raizada, M.B.; Bhardwaja, R.C. and Jain, S.K.** (1957). Grasses of the Upper Gangetic Plain. Part I (Maydeae to Andropogoneae). *Indian For. Rec. (NS) Bot.*, **4**(7) : 171-277.
- Raizada, M.B.; Jain, S.K. and Bahadur, N.** (1983). Grasses of the Upper Gangetic Plain. Part III (Poaicoideae). Jugual Kishore and Co., Dehradun.
- Randhawa, M.S.** (1983). Flowering trees. National Book Trust, New Delhi.
- Rao, R.S.** (1986a). Flora of Goa, Diu, Daman, Dadra and Nagarhaveli I and II. Flora of India, Series 2, *Bot. Sur. India, Howrah*.
- Rau, M.A.** (1969). Flora of the Upper Gangetic Plain and of the Adjacent Siwalik and Sub-Himalayan Tracts - Check list. *Bull. Bot. Sur. India. (Suppl.)* **2**: 1-87.
- Roxburgh, W.** (1820-1824). Flora indica (Edited by W. Carey and N. Wallich). Serampore, 2 Vols.
- Roxburgh, W.** (1832). Flora indica (Edited by W. Carey). Serampore, 3 Vols.
- Sabnis, T.S.** (1940-41). A contribution to the Flora of Punjab Plains and its associated hill regions. *J. Bombay Nat. Hist. Soc.*, **42**:124-149, 342-379, 533-586.
- Santapau, H.** (1950). Notes on the Scrophulariaceae of Bombay. *J. Bom. Nat. Hist. Soc.*, **49**: 25-49.
- Santapau, H.** (1953). Notes on the Acanthaceae of Bombay. *J. Bom. Nat. Hist. Soc.*, **51**: 349-368.
- Santapau, H.** (1953). The genus *Dipteracanthus* Nees in India. *Ind. For.*, **79**: 321-323.
- Santapau, H.** (1956). The botanical exploration of India. *J. Indian Bot. Soc.*, **35**: 1-6.
- Santapau, H.** (1957). *Eclipta prostrata*, *E. erecta* or *E. alba* which is correct name. *J. Bom. Nat. Hist. Soc.*, **54**: 475-476.
- Santapau, H.** (1958). Floristic studies in India. *Memoir Indian Bot. Soc.*, **1**:117-121.
- Santapau, H.** (1962). The present state of taxonomy and Floristics in India after independence. *Bull. Bot. Surv. India*, **4**:209-216.
- Saxena, H.O.** (1967) New plant records for the Upper Gangetic Plain. *Ind. For.*, **93**: 329.
- Sharma, S. and Tyagi, B.** (1979) Flora of North-East Rajasthan. Kalyani Publishers. New Delhi.
- Somdeva** (1968). A critique on Duthie's Flora of the Upper Gangetic Plain. *Bull. Bot. Sur. India*. **10**: 177-182.

- Somdeva and Naithani, H.B.** (1974). Cyperaceae of Dehradun valley and the Adjacent Siwaliks. *Ind. For.*, **100**(10): 636-654.
- Somdeva and Naithani, H.B.** (1975). Addition to the Cyperaceae of Dehradun valley and Adjacent Siwaliks. *Ind. For.*, **101**(3):205.
- Srivastava, T.N.** (1976). Flora Gorakhpurensis. Today and Tomorrow's Printers and Publishers, New Delhi.
- Stant, Margaret Y.** (1952). Anatomical evidence for including Nyctanthes and Dimetra in the Verbenaceae. *Kew Bull.*, 273-276.
- Stewart, J.L.** (1869). Punjab Plants comprising Botanical and Vernacular Names and Uses of Most of the Trees. Shrubs and Herbs of Economic Value Growing within the Province, Lahore.
- Subramanyam, K.** (1962). Aquatic Angiosperms. Botanical monograph. No. 3, CSIR. New Delhi.
- Subramanyam, K. and Sreemadhavan, C.P.** (1970). The permanent role of herbaria in modern taxonomic research. *Bull. Bot. Sur. India*, **12**: 210-212.
- Thothathri, K.** (1982). Leguminoceae : Genus-Derris. In Fascicles of Flora of India. No. 8, *Botanical Survey of India, Howrah.*
- Voss, E.G.** (1983). International code of botanical Nomenclature. Bohn, Scheltema and Holkema, Utrecht/Antwerpen, Dr. W. Junk Publishers. The Hague/Baston.
- Wight, R.** (1983-1953). Icones Plantarum Indiae orientalis. Madras. 6 Vols.



## JOURNAL OF PLANT DEVELOPMENT SCIENCES

(AN INTERNATIONAL QUARTERLY RESEARCH JOURNAL)

Editor-in-chief invites manuscripts for the publication in new esteem Journal of Plant Development Sciences. Journal of Plant Development Sciences is an international quarterly research Journal and promoted to:

- Basic & Applied Plant Sciences
- Agricultural Sciences
- Ethnobotany
- Taxonomy
- Pharmacognosy
- Phytochemistry
- Pharmacology
- Biochemistry
- Microbiology
- Biotechnology
- Environmental Sciences
- Horticulture
- Forestry
- Field Botany

Manuscripts for the publication may be submitted to the  
**Editor-in-chief, Journal of Plant Development Sciences,**  
**121/8, Pragati Nagar, Saket, Meerut – 250001 (U.P.) INDIA**

**Email-** amittomar1982@yahoo.co.in