

EFFECT OF NITROGEN AND POTASSIUM FERTIGATION SCHEDULES IN RABI SUNFLOWER ON PLANT GROWTH PARAMETERS

K. Preethika Reddy*, M. Uma Devi, V. Ramulu and M. Madhavi

Water Technology Centre, Professor Jayashankar Telangana State Agricultural University
Rajendranagar, Hyderabad 500 030
Email: preethikareddy13193@gmail.com

Received-06.04.2021, Revised-19.04.2021, Accepted-28.04.2021

Abstract: A field experiment was conducted at Water Technology Centre, College farm, Rajendranagar, Hyderabad with sunflower (variety DRS-1) during *rabi* 2017-18 in a randomized block design with three replications and the treatments were nine with combinations of N (75 kg ha⁻¹) and K (30 kg ha⁻¹) fertilizers applied by fertigation through ventury at different intervals *viz.*, 3 days and 4 days. Drip irrigation was scheduled once in 2 days at 0.8 E pan. Fertigation was imposed at 16 DAS to 88 DAS and completed in 19 and 10 splits in 4 and 8 days interval respectively. The source of N and K fertilizers was urea and potassium sulphate respectively. The soil was sandy clay loam in texture, alkaline in reaction, non saline, low in available nitrogen, medium in available phosphorus and potassium. The amount of total irrigation water applied was 3188 m³ and 4666 m³ in drip irrigation and furrow irrigation treatments, respectively. The N and K fertigation (75-30 kg N-K₂O ha⁻¹) at 4 days interval has recorded relatively higher plant growth parameters like number of leaves plant⁻¹ (25.5), plant height (208 cm), SPAD Chlorophyll Meter Reading (55.7), leaf area index (3.5) and dry matter (158.1 g m⁻²) which were on par with N and K at 8 days interval.

Keywords: Sun flower, N &K Fertigation schedule, Growth parameters, SPAD, LAI

INTRODUCTION

The projected population of India is expected to be around 1.48 billion by 2030 and the required oil seeds production will be around 102.3 million tonnes by 2030 (DRMR, Vision 2030). With the burgeoning population, improved living standard and purchasing power of the people, the demand for vegetable oil in the country is increasing at the rate of about 4-6 per cent. Thus, the country is constrained to import edible oil in large quantities involving huge expenditure in foreign exchange. It is imperative to enhance the domestic production to meet the increasing demand for edible oils. Therefore there is an urgent need to improve the productivity of oilseed crops to bridge up the current demand-supply gap. Till now there is no major breakthrough to increase the production of vegetable oil through traditional crops. On the other hand fresh water available for agriculture is decreasing due to increasing demand from industrial and home consumption sectors of water usage. Hence, a sustainable method to improve the water use efficiency and water productivity has to be evolved in arid and semiarid regions where water scarcity is already experienced. Drip irrigation is the one option available to optimize agricultural production with limiting water. It can save water up to 40 to 70% as well as increase the crop production to the extent of 20 to 100% (Reddy and Reddy, 2003). Water productivity is three times more by drip irrigation than the furrow irrigation (Kadasiddappa, 2015). Fertigation which is a sophisticated and efficient method of applying fertilizer through the irrigation system as a carrier and distributor of crop nutrients

holds the key (Bachchhav, 2005). Application of fertilizers along with irrigation water through drip fertigation can improve sunflower yield and fertilizer use efficiency and meets crop demand throughout the crop growing season. Earlier studies (Himaja, 2017) indicated the optimum NPK level as 75-90-30 kg N-P₂O₅-K₂O ha⁻¹ by fertigation for sunflower in Southern Telangana Zone. The present study was planned to identify the optimum schedule of N and K fertilizers for sunflower by fertigation.

MATERIALS AND METHODS

The experiment was conducted during *rabi* 2017-18 with variety DRS-1 at Water Technology Centre, College farm, College of Agriculture, Rajendranagar, Professor Jayashankar Telangana State Agriculture University, Hyderabad on a sandy clay loam soil, alkaline in reaction, non saline, low in available nitrogen, medium in available phosphorus and potassium. The bulk density of soil at 0-15 cm depth was 1.46 mg m⁻³ with available soil moisture 32.6 mm. The irrigation water was alkaline (pH=8.1) and categorized under the Class II (C₃S₁) suggesting that it is suitable for irrigating the crop by following good management practices. There was no carbonates hazard. The drip specifications include laterals of 16 mm diameter laid at 1.2 m apart with spacing of 0.5 m distance between two inline emitters. The emitter discharge was 4.0 lph. The experiment was laid out in a randomized block design with three replications and nine treatments at 100% RDF *viz.*, no application of N and K₂O fertilizers + drip irrigation (T₁), manual application of N and K fertilizers + drip irrigation (T₂), application of only N (75 kg ha⁻¹)

*Corresponding Author

through fertigation at 4 days interval (T_3), application of only N (75 kg ha^{-1}) through fertigation at 8 days interval (T_4), application of only K (30 kg ha^{-1}) through fertigation at 4 days interval (T_5), application of only K (30 kg ha^{-1}) through fertigation at 8 days interval (T_6), application of N and K ($75\text{ kg N} - 30\text{ kg K}_2\text{O ha}^{-1}$) through fertigation at 4 days interval (T_7), application of N and K ($75\text{ kg N} - 30\text{ kg K}_2\text{O ha}^{-1}$) through fertigation at 8 days interval (T_8) and manual application of N and K fertilizers + furrow irrigation (T_9). The recommended dose of fertilizer (RDF) 75, 90 and 30 kg N, P_2O_5 and $\text{K}_2\text{O ha}^{-1}$, respectively was applied. A common dose of phosphorus was applied to all the treatments including control (T_1). Nitrogen and potassium were applied through drip fertigation at different growth stages as per treatments through urea and potassium sulphate respectively. For manual application treatments (T_2 and T_9) N was applied in three equal splits at basal, 30 and 50 DAS through urea and potassium was applied through muriate of potash (MOP) in a single basal dose. Irrigation scheduling was done as per the treatments. Scheduling of irrigation for treatments T_1 to T_8 (except T_9) were fixed for once in two days based on daily evaporation data recorded from USWB class 'A' pan evaporimeter in agro-meteorological station, ARI Farm, Rajendranagar, Hyderabad at 0.8 Epan and furrow irrigation (T_9) was at 1.0 IW/CPE ratio with 50 mm irrigation depth in furrows in between paired rows (80 cm/40 cm). There was no rainfall during the crop growth period. The amount of water applied under drip irrigation was 318.8 mm and under surface irrigation was 466.6 mm. Fertigation was given at 4 and 8 days intervals starting from 16 DAS to 88 DAS. The fertigation schedule is detailed in Table 1. Sunflower hybrid DRSH-1 was sown by a paired row system on 16th November 2017 by adopting a spacing of 80/40 cm between the rows and 25 cm between the plants to maintain a desired plant population of 66,666 plants ha^{-1} .

RESULTS

Data on plant height is presented in Table 2. It ranged from 35.3 to 41.2 cm, 135.9 to 181.7 cm, 158.8 to 206.7 cm and 161.7 to 208.1 cm respectively at 30, 60, 90 DAS and at harvest respectively. Results revealed that plant height increased from 30 to 90 DAS rapidly and afterwards increase was at a slower pace up to harvest. It was significantly influenced by treatments at all the growth stages except at 30 DAS. At 60, 90 DAS and at harvest the highest plant height was observed under treatment T_7 (181.7 cm, 206.7 cm and 208.1 cm) which was on par with treatments T_8 (177.2 cm, 201.8 cm and 203.6 cm), T_3 (174.1 cm, 195.0 cm and 196.8 cm) and T_4 (172.6 cm, 194.9 cm and 196.7 cm). The lowest plant height was recorded under treatment T_1 (Control) (135.9 cm, 158.8 cm and 161.7 cm). The N + K at 4 days interval (T_7) has recorded higher plant height compared to manual application of N and K fertilizers + drip irrigation (T_2) and manual application of N and K fertilizers + drip irrigation (T_2) has recorded an increase in plant height compared to manual application of N and K fertilizers + furrow irrigation (T_9). Application of fertilizer especially nitrogen enhances chlorophyll synthesis resulting in increased carbohydrate assimilation, which in turn responsible for higher vegetative growth. Due to frequent application of water and fertilizers by drip fertigation, nutrients were effectively utilized due to direct contact with the root system and least loss of nutrients through leaching. Similar results for higher plant height at 100 % RDF through fertigation over manual application of 100 % RDF through surface irrigation were reported by Sanju *et al.* (2013) and Soni *et al.* (2017).

Table 1. Plant height (cm) of *rabi* sunflower as influenced by N and K fertigation schedules

Treatments	Plant height (cm)			
	30 DAS	60 DAS	90 DAS	At harvest
T_1 - Control (N_0K_0)	35.3	135.9	158.8	161.7
T_2 - Manual application of N and K + drip	40.2	162.4	187.5	188.2
T_3 - Fertigation of N at 4 days interval	37.8	174.1	195.0	196.8
T_4 - Fertigation of N at 8 days interval	37.7	172.6	194.9	196.7
T_5 - Fertigation of K at 4 days interval	37.6	147.6	180.4	182.5
T_6 - Fertigation of K at 8 days interval	37.0	145.6	171.9	173.7
T_7 - Fertigation of N and K at 4 days interval	40.0	181.7	206.7	208.1

T ₈ - Fertigation of N and K at 8 days interval	39.8	177.2	201.8	203.6
T ₉ - Manual application of N and K + furrow	41.2	148.1	169.1	170.7
SEm ±	1.91	4.72	5.93	5.65
CD (p=0.05)	NS	14.29	17.94	17.07

The no. of functional leaves plant⁻¹ at 30, 60, 90 DAS and at harvest are presented in Table 3 and it ranged from 9.9 to 11.4, 18 to 25.5 and 14.0 to 19.7 respectively. The number of functional leaves plant⁻¹ increased with crop ontogeny up to 60 DAS and thereafter started declining until harvesting due to remobilization of nutrients from source to sink and rapid senescence of leaves. The number of leaves plant⁻¹ due to different treatments was not significantly influenced at all the growth stages of

study except at 60 DAS. At 60 DAS maximum number of leaves plant⁻¹ were recorded with T₇ (N + K at 4 days interval) (25.5) which has 10.9 % more number of leaves compared to manual application of N and K fertilizers + drip irrigation (T₂) and the least was recorded under treatment T₁ (Control) (18.0). At 90 DAS, the maximum number of leaves plant⁻¹ recorded with T₇ (N + K at 4 days interval) (19.7) and the lowest was in T₉ (manual application of N and K fertilizers + furrow irrigation) (14.0).

Table 2. Number of leaves plant⁻¹ of *rabi* sunflower as influenced by N and K fertigation schedules.

Treatments	Number of leaves plant ⁻¹		
	30 DAS	60 DAS	90 DAS
T ₁ - Control (N ₀ K ₀)	9.9	18.0	17.0
T ₂ - Manual application of N and K + drip	11.4	23.0	17.3
T ₃ - Fertigation of N at 4 days interval	11.0	24.3	19.0
T ₄ - Fertigation of N at 8 days interval	10.9	24.1	18.1
T ₅ - Fertigation of K at 4 days interval	10.5	20.8	17.0
T ₆ - Fertigation of K at 8 days interval	10.3	20.1	16.9
T ₇ - Fertigation of N and K at 4 days interval	11.1	25.5	19.7
T ₈ - Fertigation of N and K at 8 days interval	11.1	24.4	19.1
T ₉ - Manual application of N and K + furrow	11.3	20.5	14.0
SEm ±	0.86	0.78	1.35
CD (p=0.05)	NS	2.38	NS

The data on spad chlorophyll meter readings (SCMR) at 30, 60, 90 DAS and at harvest are presented in Table 3 and it ranged from 30.8 to 37.1, 33.2 to 55.7, 24.7 to 44.9 and 23.2 to 39.2 respectively. The readings were not significantly influenced by different fertigation schedules at 30 DAS and significantly influenced during the other stages. The SCMR readings at 60 DAS were observed to be higher when compared to other stages of crop growth. During all the growth stages, the lowest SCMR was noticed in T₁ (Control) and the highest was noticed with T₇ (N + K at 4 days interval) which was significantly higher over application of only K at 4 and 8 days interval treatments (T₆ and T₇) and manual application of N and K treatments (T₂ and T₉) respectively. Whereas,

the SCMR recorded with T₈ (N + K at 8 days interval), T₃ (N at 4 days interval), T₄ (N at 8 days interval) treatments was found to be at par with T₇. At 60, 90 DAS and at harvest, N + K at 4 days interval (T₇) has recorded 15.8, 11.9, 13.9 % higher SCMR compared to T₂ (manual application of N and K fertilizers + drip irrigation) treatment respectively. Manual application of N and K fertilizers + drip irrigation (T₂) has recorded 18.3, 21.1, 12.0 % higher SCMR compared to manual application of N and K fertilizers + furrow irrigation (T₉). The phenomenon of increased chlorophyll content with increased N nutrition under fertigation could be due to uniform distribution and adequate availability of nutrients and water under drip fertigation in the root zone of the crop (Gupta *et al.*, 2010 and Aminifard *et al.*, 2012).

Table 3. SPAD Chlorophyll Meter Reading (SCMR) of *rabi* sunflower as influenced by N and K fertigation schedules.

Treatments	SCMR			
	30 DAS	60 DAS	90 DAS	At harvest
T ₁ - Control (N ₀ K ₀)	30.8	33.2	24.7	23.2
T ₂ - Manual application of N and K + drip	36.9	48.1	40.1	34.4
T ₃ - Fertigation of N at 4 days interval	36.5	54.1	43.5	37.9
T ₄ - Fertigation of N at 8 days interval	36.4	53.8	42.8	37.8
T ₅ - Fertigation of K at 4 days interval	36.1	40.3	32.4	30.1
T ₆ - Fertigation of K at 8 days interval	36.0	39.9	32.1	29.7
T ₇ - Fertigation of N and K at 4 days interval	37.1	55.7	44.9	39.2
T ₈ - Fertigation of N and K at 8 days interval	36.6	54.6	43.6	38.2
T ₉ - Manual application of N and K + furrow	36.4	40.6	33.1	30.7
SEm ±	1.51	1.57	1.78	1.49
CD (p=0.05)	NS	5.84	5.38	3.16

Leaf area index is defined as the ratio of green leaf area to a unit ground area, which is one of the important indicators of growth and productivity of the crops. The data is presented in Table 5 and indicated that LAI ranged from 0.5 to 0.6, 1.5 to 3.7, 1.2 to 2.5 and 0.4 to 0.8 at 30, 60 and 90 DAS and at harvest, respectively. The LAI significantly influenced by treatments at all the growth stages studied except at 30 DAS. Relatively higher LAI values were noticed at 60 DAS. At 60, 90 DAS and at harvest the lowest LAI values were noticed with T₁ (Control) and the highest was noticed in T₇ (N + K at 4 days interval). At 60, 90 DAS and at harvest N + K at 4 days interval (T₇) has recorded 42.3, 38.8, and 60 % increase in LAI compared to T₂ (manual application of N and K fertilizers + drip irrigation) treatment, respectively. At 60 and 90 DAS manual application of N and K fertilizers + drip irrigation

(T₂) has recorded 23.8 and 20.0 % increase in LAI compared to manual application of N and K fertilizers + furrow irrigation (T₉) respectively. Relative water content (RWC) is considered a measure of plant water status, reflecting the metabolic activity in tissues and used as a most meaningful index for dehydration tolerance. The periodical RWC was recorded by taking leaf fresh weight, dry weight and turgid weights. Data pertaining to RWC is presented in Table 6. The RWC of *rabi* sunflower at 30, 60 and 90 DAS was not significantly influenced by the different drip fertigation schedules and it ranged from 71.2 to 83.5%, 73.3 to 84.8% and 70.4 to 81.7% at 30, 60 and 90 DAS, respectively. At all the crop growth stages the highest RWC was observed with T₇ (N + K at 4 days interval) and lower RWC was with T₁ (control).

Table 4. Leaf area index of *rabi* sunflower as influenced by N and K fertigation schedules.

Treatments	Leaf Area Index			
	30 DAS	60 DAS	90 DAS	At harvest
T ₁ - Control (N ₀ K ₀)	0.5	1.5	1.2	0.4
T ₂ - Manual application of N and K + drip	0.6	2.6	1.8	0.5
T ₃ - Fertigation of N at 4 days interval	0.6	3.3	2.2	0.6
T ₄ - Fertigation of N at 8 days interval	0.6	3.2	2.2	0.6
T ₅ - Fertigation of K at 4 days interval	0.6	2.2	1.7	0.5
T ₆ - Fertigation of K at 8 days interval	0.6	1.7	1.6	0.5

T ₇ - Fertigation of N and K at 4 days interval	0.6	3.7	2.5	0.8
T ₈ - Fertigation of N and K at 8 days interval	0.6	3.5	2.2	0.8
T ₉ - Manual application of N and K + furrow	0.7	2.1	1.5	0.5
SEm ±	0.04	0.15	0.11	0.03
CD (p=0.05)	NS	0.46	0.35	0.10

Table 5. Relative water content (%) of *rabi* sunflower as influenced by N and K fertigation schedules.

Treatments	Relative water content (%)		
	30 DAS	60 DAS	90 DAS
T ₁ - Control (N ₀ K ₀)	71.2	73.3	70.4
T ₂ - Manual application of N and K + drip	79.0	80.9	75.7
T ₃ - Fertigation of N at 4 days interval	82.3	84.3	80.9
T ₄ - Fertigation of N at 8 days interval	81.0	81.4	79.7
T ₅ - Fertigation of K at 4 days interval	81.3	83.1	79.9
T ₆ - Fertigation of K at 8 days interval	80.0	81.8	78.8
T ₇ - Fertigation of N and K at 4 days interval	83.5	84.8	81.7
T ₈ - Fertigation of N and K at 8 days interval	81.4	83.2	78.9
T ₉ - Manual application of N and K + furrow	76.9	77.8	74.4
SEm ±	2.4	2.47	2.27
CD (p=0.05)	NS	NS	NS

The data of dry matter production (DMP) at 30, 60, 90 DAS and at harvest is presented in Table 7 and it ranged from 18.5 to 21.3 g, 30.7 to 108.8 g, 63.3 to 131.6 g plant⁻¹ and 81.8 to 158.1 g plant⁻¹, respectively. Dry matter plant⁻¹ was increased gradually up to harvest. The dry matter production was significantly influenced by different treatments at all the growth stages except 30 DAS. At 60, 90 DAS and at harvest the highest dry matter plant⁻¹ was observed in T₇ (N + K at 4 days interval) (108.8 g, 131.57 g and 158.1 g) and the lowest was recorded in control (T₁).

Significant positive correlations were noticed between the seed yield of sunflower and different

growth parameters presented in Table 8. The dependence of seed yield on plant height, number of leaves plant⁻¹, SCMR, LAI and dry matter plant⁻¹ was evident from significant (p=0.01) and positive correlation between yield and plant height, number of leaves plant⁻¹, SCMR, LAI and dry matter plant⁻¹ data. The determination coefficient (R²) was observed to be 0.54 for plant height at 90 DAS, 0.46 for number of leaves plant⁻¹ at 60 DAS, 0.57 for SCMR at harvest, 0.46 for LAI at 60 DAS and 0.68 for dry matter plant⁻¹ at 60 DAS which showed a linear increase in the growth parameters with the corresponding increase in yield.

Table 6. Dry matter production (g plant⁻¹) of *rabi* sunflower as influenced by N and K fertigation schedules.

Treatments	Dry matter production (g plant ⁻¹)				Seed yield (kg ha ⁻¹)
	30 DAS	60 DAS	90 DAS	At harvest	
T ₁ - Control (N ₀ K ₀)	18.5	30.7	63.3	81.8	1262
T ₂ - Manual application of N and K + drip	19.9	76.0	108.3	134.1	2158
T ₃ - Fertigation of N at 4 days interval	19.2	99.0	93.2	122.4	1983
T ₄ - Fertigation of N at 8 days interval	19.1	92.8	88.1	118.5	1918

T ₅ - Fertigation of K at 4 days interval	19.1	54.3	79.8	109.3	1662
T ₆ - Fertigation of K at 8 days interval	18.6	52.9	71.8	102.2	1602
T ₇ - Fertigation of N and K at 4 days interval	19.7	108.8	131.6	158.1	2623
T ₈ - Fertigation of N and K at 8 days interval	19.5	99.9	126.2	146.9	2437
T ₉ - Manual application of N and K + furrow	21.3	59.1	89.3	110.1	1795
SEm ±	0.93	5.30	6.04	6.41	98.90
CD (p=0.05)	NS	16.01	18.49	23.80	299.07

Table 7. Correlation and regression between seed yield Vs growth parameters of *rabi* sunflower as influenced by N and K fertigation schedules.

S.No.	Parameter	DAS	r ² value	Regression Equation	R ²
1.	Plant height (cm)	30	0.401*	y = 52.32x - 78.85	0.17
		60	0.717**	y = 17.93x - 946.3	0.51
		90	0.734**	y = 18.69x - 1516	0.54
		Harvest	0.694**	y = 17.6x - 1355	0.48
2.	Number of leaves plant ⁻¹	60	0.682**	y = 147.9x - 1537	0.46
3	SCMR	30	0.484*	y = 92.52x - 1422	0.23
		60	0.664**	y = 47.91x - 279	0.44
		90	0.683**	y = 50.86x - 18.20	0.47
		Harvest	0.758**	y = 68.71x - 381.6	0.57
4.	LAI	60	0.677**	y = 563.2x + 580.4	0.46
		90	0.594**	y = 943.4x + 280.7	0.35
		Harvest	0.656**	y = 2217x + 625.5	0.43
5.	Dry matter (g plant ⁻¹)	60	0.828**	y = 14.26x + 868.3	0.68
		90	0.780**	y = 15.37x + 489.4	0.61
		Harvest	0.765**	y = 13.60x + 257.8	0.59

CONCLUSION

Based on the results obtained in the present investigation, it can be concluded that the sunflower crop grown with drip fertigation at 100 % RD of N + K (75 kg N - 30 kg K₂O ha⁻¹) with 4 days interval during *rabi* under Hyderabad semi arid conditions recorded the highest plant growth parameters viz., plant height (203.6 cm), number of leaves plant⁻¹ (25.5), SCMR (55.7), LAI (3.7) and dry matter

production (158.1 g plant⁻¹) and can be recommended for application.

REFERENCES

Aminifard, M.H.; Arojee, H.; Ameri, A. and Fatemi, H. (2012). Effect of plant density and nitrogen fertilizer on growth, yield and fruit quality of sweet pepper (*Capsicum annum* L.). *Asian Journal of Agricultural Research*, 7(6): 859-866.

- Bachchhav, S.M.** (2005). Role of fertigation in improving fertilizer use efficiency. *Fertilizer Marketing News*, 36: 1-9.
- DRMR, Vision.** (2030). Directorate of Rapeseed-Mustard Research, Bharatpur, Rajasthan
- Gupta, A. J.; Ahmed, N.; Bhat, F. N. and Chattoo, M. A.** (2010). Production of hybrid tomato for higher income under drip irrigation and fertigation in Kashmir valley. *Indian Journal of Horticulture*, 67(1): 127-131.
- Kadasiddappa, M.; Praveen Rao, V.; Yella Reddy, K.; Ramulu, V.; Uma Devi, M.; Narender Reddy, S. and Mallappa, B.V.** (2015). Effect of drip and surface furrow irrigation on growth, yield and water use efficiency of sunflower. *An International Journal*, 10(7): 3872-3875.
- Reddy and Reddy** (2003). Principles of Agronomy. Kalyani Publication, New Delhi, India. pp. 319-320.
- Sanju, H.R.** (2013). Effect of precision water and nutrient management with different sources and levels of fertilizer on yield of groundnut. *M.Sc Thesis*. University of Agricultural Sciences, GKVK, Bangalore, India.
- Soni, J.K.; Raja, N.A. and Vimal, K.** (2017). Study of comparative performance of drip, micro sprinkler fertigation system and surface irrigation in groundnut (*Arachis hypogaea* L.). *Environment and Ecology*, 35(3): 050-2055.

