

# SEED PROTEIN PROFILING THROUGH ELECTROPHORESIS IN MUNGBEAN [*VIGNA RADIATA* (L.) WILCZEK]

Subek Singh, Prafull Kumar<sup>1</sup> Ankita Trivedi and Avadhesh Kumar Koshal\*<sup>2</sup>

Department of Genetics and Plant breeding, C.S. Azad University of Agriculture and Technology,  
Kanpur-208 002, (Uttar Pradesh) India

<sup>1</sup>Sardar Vallabhbhai Patel University of Agriculture & Technology,  
Modipuram, Meerut, (Uttar Pradesh) India

<sup>2</sup>ICAR- Project Directorate for Farming Systems Research,  
Modipuram, Meerut, (Uttar Pradesh) India

Received-05.04.2015, Revised-22.04.2015

**Abstract:** Mungbean (*Vigna radiata* (L.) Wilczek) is an important pulse crop in India with advancement and development of hundreds of varieties and introduction of intellectual property rights it is necessary to identifying them individually for identification and registration purposes. The present investigation was carried out during 2012-2013 in biotechnology lab, Department of genetics & Plant Breeding, C.S. Azad university of agriculture and technology, Kanpur with 13 genotypes of Mungbean K851, SML-683, KM 2272, PDM 51, T- 44, KM-2260, HUM-12, HUM-6, Pusa Vishal, IPM 02-3, IPM 2-14, KM-2241, PDM-139, for protein profiling through SDS-PAGE. In present investigation, 13 variety of Mungbean were studied for varietal identification through electrophoresis. Protein was extracted from dry seed of mungbean varieties and analysed by SDS-PAGE. On the basis of photographs, electrophoregrams, Rm values and dendograms (UPGMA cluster analysis) of banding patterns through SDS-PAGE, results found that the number of protein bands found in 13 genotypes ranged from 10 to 17 with Rm value 0.08 to 0.97 for tris soluble proteins. Protein banding pattern of tris soluble proteins was found more distinct in SDS-PAGE. In UPGMA cluster analysis all the genotypes fall in seven cluster groups. SDS-PAGE for tris soluble proteins found suitable for testing distinctness, uniformity, stability of varieties for registration and identification. On the basis of results, this can be said for characterization and identification of genotypes of mungbean, that electrophoretic profile for tris soluble proteins through SDS-PAGE was resulted distinct banding pattern and act as 'genotypic finger printing'. Therefore, electrophoregram of tris soluble protein in SDS-PAGE was found much better for identification of genotypes in mungbean.

**Keywords:** Mungbean, SDS-PAGE, Varietal identification, UPGMA

## INTRODUCTION

Mungbean (*Vigna radiata* Wilczek) is short duration legume crop belong to the order leguminosae and papilionoideae family. It is a self-pollinated diploid legume with the chromosome number  $2n= 22$  the mungbean crop having high nutritive value 100 g of mungbean contain 23.83 g protien. most of which is consumed in the domestic market with the advancement and development of thousands of improved new varieties and introduction of IPR, it is necessary to identifying them individually for identifications and registration purposes. For the purpose necessity of quick, reliable and reproducible laboratory techniques are required. Protein profiling through SDS-PAGE is an alternate techniques for distinguishing the genotypes. Protein markers are stable, reproducible and genetically controlled and can be conducted in relatively short time.the protein profiling of seed storage proteins in cultivated lentil and their significantly differences in banding patterns by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE). In electrophoresis, substances in a mixture, which are ionisable, can be reported from others that are net, by subjecting the mixture to an electric field:

SDS-PAGE is the most commonly methods used for studying protein differences between species, previously classified on the basis of their morphological characters. SDS is an anionic detergent which binds strongly and denatures proteins. The number of SDS molecule bound to a polypeptide chain is approximately half the no. of amino acid residues in that chain. The protein SDS complex carries net negative charges. Hence, more towards the anode and the separation is based on the size of the protein.

## MATERIAL AND METHOD

The study was conducted during 2012-13 at Biotechnology Lab, Department of Genetics and Plant breeding, C.S. Azad University of Agriculture and Technology, Kanpur (UP). For the study, genetically pure seeds of 13 mungbean genotypes K857, SML-683, KM 2272, PDM 51, T- 44, KM-2260, HUM-12, HUM-6, Pusa Vishal, IPM 02-3, IPM 2-14, KM-2241, PDM-139 were procured from the mungbean Breeder of the university on the basis of morphological characters. The soluble proteins were analysed through SDS-PAGE method recommended by Dadlani *et al.*, 1993 for variety identification

\*Corresponding Author

### Preparation of Sample

About 1 g seed were grinded in mortar and pestle after removing the seed coat and defatted by defatting solution 4 times of all 13 genotypes were taken in tube separately 1 ml Tris-glycine extraction buffer (pH 8.3) was added to 0.5g of defatted powder and left over night. 10 % solution of SDS (10 $\mu$ l), 2-mercapto ethanol (10 $\mu$ l) with bromophenol blue (10  $\mu$ l) was added. Mixed well and left over night in a refrigerator. The sample was heated in boiling water bath kept for 10 minutes in water bath for 10 minutes at 100°C. . The tubes were cooled and centrifuged at 10000 rpm for 10 minutes. The clear supernatant was used for electrophoresis.

### Preparation of gel

Seed protein were analysed through slab type SDS-PAGE followed by Laemmli (1970) using 12% polyacrylamide gel. Electrophoresis was conducted in Atto Electrophoresis Unit using fourteen well for loading the sample. fixed the gel cassette into the electrophoresis unit as per design of the equipment. Loaded 50  $\mu$ l of clear supernatant with one drop of tracking dye (Bromophenol blue) was loaded to each well. The electrophoresis was conducted at a constant current of 1.5 mA per well [42 mA] till the tracking dye crossed the stacking gel than current was fixed @ 2 mA per well at 220 V. The electrophoresis was stopped after the tracking dye reached the bottom of the gel.

### Fixing and staining

Removed the cassette from the unit and take out the gel gently. Placed it in a staining tray and incubate overnight in 15% Trichloroacetic acid solution. Wash thoroughly the excess SDS, which might precipitate on the surface. Sufficient 1% comassie brilliant blue solution, prepared in methanol was added to cover the gel uniformly and incubated for

16 hr to stain than rinsed with water. Destaining in water and 5% Acetic acid for two days clears the gel background, resulting in a better resolution. The gel was placed over a trans-illuminator to draw the electrophoregram for calculating Rm values.

### RESULT AND DISCUSSION

Results obtained through SDS-PAGE showed that the method provided a powerful tool for reliable variety discrimination and identification based on genetic differences of seed storage protein composition. Genotypes were distinguished on the basis of presence and absence of protein bands at particular Rm value and total numbers of bands present.

In the electrophoregram of tris soluble protein through SDS-PAGE (Fig. 1 & 2) in 13 genotypes, the numbers of protein bands were ranged from 10 to 17 with Rm value 0.08 to 0.97.

The mungbean genotypes based on similarity distance dendrogram of 13 genotype of tris soluble protein banding pattern using UPGMA clusters analysis (Fig. 3) were grouped in 8 clusters. Clusters first contain three genotypes in grouped namely ; K-851, KM-2260 and SML-683, in which K-851 and KM-2260 are more close than SML-683. cluster, second and third KM-2272 and PDM-51, respectively have wider distance to other genotypes. Cluster fourth, IPM 02-3 and PDM-139 are grouped that are contain two genotypes which are close to each other. Cluster fifth contains two genotypes T- 44 and KM- 2241 are grouped that are close to each other. Cluster six contains two genotypes Pusa Vishal, and IPM-02-14 are grouped that are close each other. Cluster seven and eight HUM-6 and HUM- 12 HUM-6, respectively have wider distance to other genotypes.





Fig.: SDS-PAGE Electrophoregrams of Tris soluble proteins in Mungbean genotype

**Table 1.** Relative mobility Rm values and presence/absence of total soluble seed protein in mungbean genotype (SDS-PAGE)

Band No.	Genotype Ram value	K 851 (1)	SML 683 (2)	KM 2272 (3)	PDM 51 (4)	T 44 (5)	KM 2260 (6)	HUM 12 (7)	HUM 6 (8)	Pasa Vishal (9)	IPM 02-3 (10)	IPM 2-14 (11)	KM 2241 (12)	PDM 139 (13)
1.	0.08	-	+	-	+	-	-	+	-	-	-	-	+	-
2.	0.11	+	+	+	+	-	-	-	+	-	+	-	+	+
3.	0.16	+	+	+	+	-	+	-	-	-	+	-	-	+
4.	0.18	-	-	-	-	+	-	+	+	-	+	-	+	+
5.	0.21	+	+	+	+	-	+	+	+	+	-	+	-	-
6.	0.24	+	+	+	-	+	+	-	-	+	-	-	+	-
7.	0.27	-	-	-	+	-	-	-	+	-	-	-	-	-
8.	0.29	+	-	-	-	+	+	+	-	-	+	+	+	-
9.	0.31	+	+	-	+	+	+	-	+	+	-	-	+	-
10.	0.36	+	+	+	-	-	+	-	+	-	+	+	+	+
11.	0.40	+	+	+	+	+	+	+	-	-	+	-	+	+
12.	0.44	-	-	-	-	-	-	+	-	-	-	-	-	+
13.	0.47	-	+	+	-	-	+	-	-	+	-	+	+	-
14.	0.51	+	-	+	+	+	+	-	+	+	+	+	-	+
15.	0.55	+	+	+	+	+	+	-	+	-	+	+	+	+
16.	0.58	+	-	+	-	+	+	+	+	+	+	-	+	-
17.	0.63	+	+	+	+	+	+	+	+	-	-	+	+	+
18.	0.70	+	+	+	+	-	+	+	+	+	+	-	+	+
19.	0.74	-	-	+	-	-	-	-	+	-	-	-	-	-
20.	0.76	+	+	-	+	+	+	+	-	+	+	+	+	-
21.	0.85	+	+	+	+	+	+	+	+	+	+	+	+	+
22.	0.90	+	+	+	+	+	+	-	-	+	+	+	+	+
23.	0.97	+	+	+	+	+	+	+	+	+	+	+	+	+
	Total	17	16	16	15	13	17	12	14	10	14	11	17	13



On the basis of photographs, electrophoregrams, Rm values and dendograms (UPGMA cluster analysis) of banding patterns through SDS-PAGE, variations

were found in Rm value of protein bands, numbers of protein bands, similarity distance cluster analysis. These results are supported to the findings of and

Kawas, A. and S. Nehir, (1992) in mungbean and lentil. Amjad *et al.*, (2012) in mungbean. Gunarti *et al* and Lee *et al* (2013) in mungbean. Protein banding pattern of tris soluble proteins were found more distinct. On the basis of results, this can be said for characterization and identification of genotypes of mungbean, that electrophoretic profile for tris soluble proteins through SDS-PAGE was resulted distinct banding pattern and much better.

## REFERENCES

- Afzal, M., Kawase, M., Nakayama, H. and Okuno, K.** (1994). Variation in electrophoregrams of total seed protein and wx protein in *foxtail millet*. *Breed Sci.*, 44: 642
- Amjad, Hameed, Tariq Mohamed Shah, Baban Manzoor Atta, Nayyar Iqbal, M., Ahsanul Haq and Hina Ali** (2009). Comparative seed storage protein profiling of kabuli chickpea genotypes. *Pakistan J.Bot.*, 41 (2): 703-710.
- Chauhan, P., Ram, C., Mann, A. and Sangwan, V.P.** (2002). Molecular weight analysis of seed proteins of forage sorghum (*sorghum bicolor* (L.) Moench). *Seed Science and technology*, 30 (1): 11-16.
- Drawn S. Luthe** (2007). Electrophoretic analysis of seed proteins in the dicotyledoneae. *Plant Molecular Biology Reporter*, 10 (3): 254-262.
- Gupta, S.K., Bala, A., Adarsh Bala and Satija, D.R.** (2002). Genetic diversity in relation to protein and protein fractions in chickpea (*Cicer arietinum*), **29** (2): 122-135
- Horax, R., Hettiarachchy, N.S., Chen, P. and Jalauddin, M.** (2006). Preparation and characterization of protein isolate from compeer (*Vigna unguiculeta* L. Walp). *Journal of Food Science*, 69 (2): 114-118.
- Iqbal, S.M., Ghaffor, A. and Ayub, N.** (2005). Relationship between SDS-PAGE markers and Ascochyta blight in chickpea. *Pak. J. Bot.*, 37: 87-96.
- Omid Sofalian and Mostafa Valizeh** (2009). Investigation of seed storage proteins in some wild wheat progenitors using SDS-PAGE and acid PAGE. *Not. Hort. Agrobot. Cluj.*, 37 (1) : 179-182
- Prasad, R., Prasad, D. and Taneja, A.D.** (2001). SDS-PAGE of soluble seed proteins for identification of cotton cultivars. *Journal of the Indian Society for Cotton Improvement*, 126: 2
- Ritu Mahajan, Sarla, P. Malhotra and Randhir Singh** (1987). Charaterization of seed proteins of urdbean (*Vigna mungo*). Springerlink Journal Article, 10: 163-173.
- Rico Juan, Julio Pster, Manual Alaiz and Javier Vioque** (2007). Electrophoretic characterization of *Ameranthus* L. Seed proteins and its systematic implications. *Botanical Journal of the Linnean Society*, 155 (1): 57-63.
- Shathe, S.K. and Salumkhe, D.K.** (2006). Solubilization and electrophoretic characterization of the great northern bean (*Phaseolas vulgaris* L.) proteins. *Journal of food science*, 46: 82.