

PERSPECTIVES ON ENHANCING VALUE OF AGRONOMIC BIOFORTIFICATION IN MAIZE

Augustine R.* and D. Kalyanasundaram

*Department of Agronomy, Faculty of Agriculture, Annamalai University,
Chidambaram, Tamilnadu, India*

Email: augustinerajendran@gmail.com

Received-07.07.2020, Revised-28.07.2020

Abstract: Most of the health components based foods are boosted by the application of mineral nutrients. Farmers fertilize the crop for optimum to higher yields which has become essential, since post green revolution. In addition to higher yield, plant nutrition also affects other human nutritional needs like proteins, oils, vitamins and minerals. Trace elements necessary to human nutrition can be optimized by applying micronutrients to food crops. Some nutrients have their own restrictions to various factors like temperature, climate, time of applications, crop adaptability, etc., and few micronutrients are beneficial and play a significant role in food nutrition making easier access in the plant edible parts by its applications. It is important to note that foliar application of Zn and Fe at the later crop stage (mid booting stage or early milking stage) is found to be effective than early applications. When compared to related interventions like supplementation and fortification, biofortification was found to be significantly cost effective in applications to crop and creates value for human nutrition.

Keywords: Iron (Fe), Zinc (Zn), Micronutrients, Supplementation, Fortification, Bio fortification

INTRODUCTION

Maize occupies third most important position in food crops next to wheat and rice. India is among the top 10 maize producers in the world and contributes around 2-3% of the total global production and is among the top 5 maize exporters in the world contributing 14% of the total maize exported to various countries around the world. Maize is an important cereal crop of India for larger section of populations, raw material for industries, feed for animals, and plays a major role in agricultural economy.

Maize is consumed as basic food in some of the major countries as, corn flakes, corn syrup and oil. For animal feed, its feed value from grain, leaf and stem is gaining more importance in many countries. For industrial and pharmaceutical applications, it can be used to produce starch, ethanol, plastics and as a base product for antibiotic production. Starch is the main product of maize which gives dextrin, liquid glucose, powder glucose and crystalline dextrose.

In the recent decades, because of its importance maize area has been steadily increasing or remaining stable in the maize producing countries. Maize processing industries encourages farmers to enhance the cultivable area under maize in order to sell the maize as cash crop. Maize is an exhaustive crop having high potential than other cereals and also absorbs huge quantity of nutrients from the soil at different growth stages of crop. The biggest user of maize in India is the poultry industry with 47% of the share followed by direct consumption at 20%. Other usages include cattle feed (14%) and starch (14%) followed by the food and beverage industry with a 7% share.

Normal maize has poor nutritional value because of poor contents of essential amino acids, such as lysine and tryptophan. But quality protein maize contains higher amount of these amino acids in the endosperm than normal maize. The productivity of quality protein maize is low due to inherent low soil fertility and poor nutrient management practices, and no use of secondary and micronutrients. The integrated use of organic manure and chemical fertilizers can augment the nutrient use efficiency and also enhance the productivity of quality protein maize. The balance combination of amino acids in quality protein maize have resulted in to its higher biological value ensuring more availability of protein to both human and animal as compared to normal maize.

Biofortification is one such unique practice done through fertilization with nutrients and is been extensively used in maize with supplemental foliar spray for increasing the high concentrations of nutrients in grain. Eventhough this practice is common in crops, spraying minerals at the appropriate time during plant growth turns to an efficient nutritional starter/gainer in plant parts which helps in direct nutritional support to human population and animals, etc.

Why biofortification?

Billions of deprived people worldwide suffer from varied physical disabilities due to (micronutrient malnutrition) insufficient supply of nutrients, amino acids, vitamins and minerals. India is also one such country where a huge malnourished people are identified which needs quick interventions. The problem of micronutrient deficiencies arises in developing countries where the rural population are totally dependent on cereal based diet as staple food. Essential human nutrition is protein, lysine, tryptophan, iron, zinc, vitamin A and vitamin C and

*Corresponding Author

their deficiency will lead to various health disorders. Similarly antinutritional compounds such as erucic acid, glucosinates and kunits trypsin inhibitor (KTI) consumed at high concentration levels will lead to complication in health of humans and livestock (Pfeiffer and McClafferty, 2007)

To overcome the nutritional problem, the most cost-effective and sustainable technique is by enriching nutrients in staple food through Biofortification. Biofortification plays a crucial role in reducing the gap between the micronutrient requirements and increased intake of three micronutrient dietaries viz., Vitamin A, Iron and Zinc which is of public health significance worldwide. The Biofortified crops are delivered with enriched nutrients to rural communities or to the lower income group, where commercial fortification or supplementation is not accessible. The primary targets of Biofortification are the women and children, where their requirements are high and this often goes as unmet.

The challenge remains, is to deliver nutritious, safe and cost-effective food to help improve the impact of nutritional security. To alleviate micronutrient deficiencies, various interventions like supplementation diet and dietary diversification has been tried and to be specific none of these approaches were found to be quiet effective, because of its poor distribution system and infrastructure and expensive (Tanumihardjo et al., 2007).

Plants are capable of synthesizing essential dietary micronutrients and act as a biochemical storage house. The plant based foods consumed abundantly from rice, wheat, cassava and maize by populations contains several micronutrients that are insufficient to meet daily nutritional requirements. These nutrients are unevenly distributed in plant parts. For example, iron and provitamin A carotenoides content is high in rice leaves but low in polished grain and similarly high in wheat bran and low in polished wheat flour. Biofortification efforts are directed towards increasing levels of micronutrients in edible tissues of crops by integrating crop management practices and genetics and breeding approaches.

Role of biofortification through fertilization

The effective role of fertilizers is to provide nutrients to plants in order to grow, mainly with Nitrogen, Phosphorus and Potassium. This approach has an advantage, because it works faster. But for a long-term strategy focussing on improving human health, there found to be a serious limitation with enhanced fertilizers. They are costly and expensive and need to be applied continuously. The potential of biofortified crops is to provide a continuous supply of micronutrients to large populations with cost effective and easily accessible (Meenakashi et al., 2010; Stein et al., 2006). With technical developments combined with economic savings Biofortification focuses on the path of “nutritional security” (Shetty, 2009). The importance of Biofortification to be noted is:

Priority in selective nutrition: Biofortification allows selected nutrition to a particular crop, either by genetic or breeding or agronomic approach. This method of approach reduces the drudgery of consuming multiple foods so that required nutrients are incorporated in the crop. Ex., Fortification of Golden rice with vitamin A, so that poor people can consume nutritious food and Orange sweet potato for Vitamin A which yielded significant results in mothers and children’s.

Need for overspending on food: In developing poor countries, people could not afford to spend on nutritious food, for which Biofortification gives a rescue hand. Since nutrition is focussed on single source, people need not spend on multiple foods. Lots of money can be saved and been used for their welfare activities.

Increases human productivity: Due to lack of nutrients, there is a reduction in efficiency in human workforce output. Substitution with nutrients in an efficient manner can improve the productivity which contributes for economic and social growth and development.

Biofortification is a multistage process which requires evaluation at every step and finally judged by the people/populations. Biofortification has many advantages on nutritional interventions. For ex., In heart disease, researchers must wait for decades to conclude if diet brings any difference. So they decided to measure the intermediate points not the final outcomes. Heath consequences of malnutrition emerges quickly, like anaemia in women, blindness in children and early death. Only biofortified food crops can make a difference and these developments and improvements will appear quickly in the coming years.

Biofortification through foliar fertilization is a new technique extensively used in maize because of the introduction of hybrids with high increased use of fertilizers containing high concentrations of nutrients with the target levels of micronutrients. Keeping in view the growing importance of maize as food and feed, Biofortification of maize includes enrichment of micronutrients like provitamin A, Fe and Zn in grain, which plays a significant role in human health system.

Micronutrient in human nutrition

Micronutrient deficiency related health issues or problems are more prevalent in both urban and rural areas. Under present scenario three billion people around the globe are deficient in vitamins, minerals, specifically vitamin A, Iodine (I), iron (Fe) and Zinc (Zn) (Dahiya et al., 2008). This is due to lack of awareness about dietary practices, poverty, and lack of affordability towards balanced foods, which leads to micronutrient deficiency.

Vitamin A: Plays an essential role in vision, immune response, bone growth, reproduction, embryonic development and regulation of adult genes. An early symptom of Vitamin A deficiency is night blindness.

Around 2.5 – 5.0 lakh children becoming blind every year and half of which die within a year of losing sight. Since humans and animals cannot synthesize vitamin A in their body, the only source achieved is through dietary means. In rural poor populations, plant foods are generally preferred than animal products, because of its rich source of provitamin A carotenoids. As a vitamin A supplement, β - Carotene rich maize is found efficient when consumed as a staple food when compared to other commercial fortified products (Gannon et al., 2014).

Iodine: It is a component of thyroid hormones. Iodine deficiency disorders leads to greater cause of preventable brain damage in the foetus and infants and retarded psychomotor development in young children. It is estimated that more than one billion individuals suffer from goiter worldwide, out of which half of them from Asia are worst affected.

Iron: An active constituent of catalytic site of heme and non-heme proteins. More than one-third population suffers from anemia out of which half of is by iron-deficiency. Iron deficiency affects cognitive development, resistance to infection, slows work productivity, dizziness and pregnancy, etc. Children of anemic mothers have low iron reserves and require more iron than supplied by breast feeding. It is estimated around 8 lakh deaths annually are attributed to iron deficiency anemia.

Zinc: It is involved in RNA and DNA synthesis and is a constituent of zinc containing enzymes mainly for its cellular growth. One third of the population is at high risk due to zinc deficiencies. Zinc deficiency leads to impaired growth, immune dysfunction, increased mortality and abnormal neuro behavioral development. Zinc deficiency is directly related to the severity and frequency of diarrhea in children causing even death.

In contrast Fe and Zn is an important nutritional problem specifically in developing countries, which needs genetic and agronomic Biofortification of food crops by enriching with Fe and Zn (other micronutrients) in grains or plant parts. Although it's very simple and cost less, the application of essential micronutrients is complicated due to method of application, soil fertility, and mineral mobility in the plant and its accumulation in different sites (Zhu et

al., 2007). This strategy has been successful in limited levels within specific locations only. Increasing the concentration of micronutrients in staple crop is the first step in making rich source of nutrition for human.

Agronomic biofortification of maize

Agronomic biofortification can provide temporary increase in micronutrient through fertilization enhancement in productive parts. Maize is one of the major staple crops with variety of uses. Maize owes worldwide significance due to its utilization in human food and livestock feed. Use of maize as human food and animal feed, has led to the increased productivity and rise in fertilizer applications and thus enhances the soil depletion and loss of phyto-availability mineral (Raut et al., 2010; Lal, 2009). Due to this, mineral deficiencies have become a critical (limiting) factor in productivity and quality of crop.

Among the Biofortification methods, application of fortified fertilizers with micronutrients is the cost effective and simplest method, and the success of agronomical Biofortification is highly variable due to its mineral mobility, its accumulation and soil compositions in the specific locations. Soil composition analysis has indicated that 1/2 of the agricultural soils are Zn deficient. Agronomic Biofortification is cost-effective and laborious as it demands continuous micronutrient application to the soils or plants. Biofortification by Fe foliar spray in rice crops was an effective way to improve iron concentration in rice grains (He et al., 2013; Yuan et al., 2013; Fang et al., 2008).

Micronutrient fertilizers should be applied regularly and are costly and environmentally biased. Overall such strategies are applied to specific crops with mineral potentiality to boost the nutritional quality of foods. In general staple crop, grains have very low bio available levels of Fe and Zn (i.e. about 5% of total Fe and 25% of total Zinc). Increasing the Fe from 5% to 20% in grains is equal to four fold increase of Fe in applications. Mainly the presence of antinutrients in plant foods plays a significant role in reducing the bioavailability of essential micronutrients (Graham et al., 2001). They are listed as:

Antinutrients	Essential micronutrients reserved
Phytic acid	Fe, Zn, Cu, Ni
Cellulose, hemicelluloses, lignin, cutin, suberin	Fe, Zn, Cu
Tannins and Polyphenols	Fe
Lectins	Fe
Goitrogens	I
Heavy Metals (e.g., Cd, Hg, Pb)	Fe, Zn

Further, it is not always possible to target the edible portion of plant parts like seed or fruit. It even results in the accumulation of nutrients in leaves or non-edible portions of plants. Therefore this technique is successful with certain minerals under specific

crops/species. In addition, fertilizer accumulation in soil and water poses environmental adverse effects. Agronomic Biofortification which includes fertilization can boost the Fe, Zn, I and Se levels in foods. Deficiencies of I and Se does not limit the

crop growth, whereas correction of Fe and Zn deficiency can benefit both crops and consumers. Timing of foliar application of micronutrients is found to be critical besides following agronomic practices in maximizing micronutrient accumulation of Zn and Fe. Application of Zn and Fe is a complimentary solution to plant breeding and both are required for better efficacies.

Importance of zinc and Fe (mineral fertilization) in maize crop

Iron (Fe) and zinc (Zn) are the two major heavy metal micronutrients which play a vital role in plants and are essential in the biological system. Increase in Nitrogen nutritional status in plants by higher N-fertilizer applications leads to higher accumulation of Fe and Zn in grain (Cakmak, 2010; white and Broadley, 2011). Although present in soil, their availability to plants is limited due to complicated soil properties and compositions (Wiedenhoeft, 2006; Cakmak, 2008). Thus Fe and Zn application is utmost important and in specific foliar application helps in absorption and control of deficiency. This found to be a best method for biofortification in maize crop.

Iron is transported around the plant in a chelated form, mainly as citrate and malate in xylem and nicotianamine (NA) synthase and its derivatives in phloem. In plants iron is stored in the form of ferritin. Plant ferritin gene has been over expressed in wheat, rice and maize, using endosperm as specific promoters. As stated earlier, bioavailability of iron is low (<15%) in plant foods and thus an important factor need to be considered in biofortification approach in staple crops like maize. Hence targeting iron specifically in endosperm is an alternative solution to the antinutrient (phytate) problem (Connorton et al., 2017).

When applied in excess, Fe fertilization remains complicated and has a strong reaction towards insolubility. A foliar application of 2-3% Fe concentration in Fe-deficient soils increases Fe level in crops and improves yield, when applied at different intervals. These are expensive and improper applications leads to environmental hazards. Iron deficiency affects more than two billion people, which directs to consider (Fe) mineral as priority in maize biofortification. Iron has low mobility in soil and due to this, foliar application is recommended with FeSO_4 and the Fe-EDDHA chelates and Fe-EDTA for better absorption (Frossand et al., 2000; Zhu et al., 2007).

The ideal Fe, provided in chelate form is more available and effective against Fe deficiency, which is quite expensive and found to be marginally better in increasing Fe concentration in grain. Foliar application of ferrous sulphate (Fe_2SO_4)₃ has shown a better effect than soil application in increasing Fe concentration in cereals (grain), which also enhances the yield of crops. Fe when used under process fortification has been successful for Fe fortification

which includes rice, fish, soy sauce, wheat flour & maize flour, milk and infant formulas.

Zinc (Zn) is the most abundant element next to iron and is an essential micronutrient for plants. The fertilization of crop with Zn either through soil or foliar application is a common practice in agriculture crop/soil where Zn deficiency plays a critical role in crop productivity. The main focus is to correct Zn deficiency and improve the yield with no attention paid from the human nutritional perspective towards the Zn concentration present in the edible parts of seeds or grains or starchy roots. Foliar application of zinc fertilizer in wheat grain increases grain zinc concentration by up to 20 ppm in India and Pakistan (Zhou et al., 2012). Usually soil applications of Zn had little effect on the Zn concentration in grain, whereas foliar Zn sprays was quiet effective in Zn accumulation in grains.

The foliar application of Zn was effective against reducing phytate (antinutrient) concentration in grain (Erdal et al., 2002; Cakmak et al., 2010). The bioavailability of Zn in diets is promising based on phytate/Zn molar ratio. In general, foliar Zn applications are Zinc Sulphate (ZnSO_4) and EDTA – Chelated Zn. For correcting Zn-deficiency, Chelated Zn is superior to ZnSO_4 but cost effective option should be compared with high priced Zn-EDTA. Above all, timing of foliar Zn-application is more important to determine the effectiveness in biofortification (Welch et al., 2013). Agronomic efficiency of Zn as foliar application was four times higher than soil applications and also rate of application was lower when applied as foliar spray (Dhaliwal et al., 2010). The dosage rate of 25 to 50 kg ZnSO_4 per ha are generally proposed for use in fertilization of soils/crops with Zn deficiency (Cakmak, 2008). Depending on the severity of the Zinc deficiency problem, a cost effective model in fertilization need to be worked out, to understand the better practices adopted in the region. At the initial stage, farmers need to be motivated about the strategy of Agronomic Biofortification use in staple crops, quality seeds and awareness about the Zinc nutrition to human health. This will bring a potential growth in fertilization (biofortification) of Zn as foliar spray or foliar + soil application, irrespective of stages it reaches for maintaining yield and helps in accumulation of available Zn in grain.

Thus enriched micronutrient (Fe and Zn) in maize overcomes malnutrition, which holds immense quality for sustainable and cost-effective solutions.

CONCLUSION

Agronomic biofortification is the simplest and fastest way for biofortification of grains with Fe, Zn or other micronutrients, where cereals are mainly the staple food in developing countries. Agronomic biofortification is an alternate route to reach the rural poor populations, who cannot afford to buy mineral

supplements nor can afford to improve the diet components. In the biofortification point of view, foliar application was found to yield better results than soil applications and also requires less amount of Fe and Zn fertilizers during applications. Adequate Fe and Zn fertilization is necessary when GM cultivars with improved Fe and Zn are developed. Thus genetic and agronomic approaches are complementary to each other and should progress in tandem. However a better understanding on micronutrient reactions, their absorption and translocation in plants, specifically to grains is required.

ACKNOWLEDGEMENT

The authors gracefully acknowledge the advice and guidance provided by Dr M. Ganapathy, Professor and Head, Department of Agronomy, Faculty of Agriculture, Annamalai University, Chidambaram, Tamilnadu, India.

REFERENCES

- Cakmak, I.** (2008). Enrichment of cereal grains with zinc: Agronomic or genetic biofortification? *Plant and Soil*, 302: 1-17.
- Cakmak, I., Pfeiffer, W.H. and McClafferty, Y B.** (2010). Biofortification of durum wheat with zinc and iron. *Cereal Chemistry*, 87: 10-20.
- Connorton, J.M., Jones, E.R., Rodriguez-Ramiro, I., Fairweather-Tait, S., Uauy, C. and Balk, J.** (2017). Wheat vacuolar iron transporter TaVIT2 transports Fe and Mn and is effective for biofortification. *Plant Physiology*, 174: 2434-2444.
- Dahiya, S., Chaudhary, D., Jaiwal, R., Dhankher, O., Singh, R.** (2008). Elemental biofortification of crop plants. In: Jaiswal P, Singh R, Dhankar OP (eds) *Plant membrane and vacuolar transporters*. CABI International, Wallingford/Cambridge, pp. 345-371.
- Dhaliwal, S.S., Sadana, U.S., Khurana, M.P.S., Dhadli, H.S. and Manchanda, J.S.,** (2010). Enrichment of rice grains through ferti-fortification. *Indian Journal of Fertilizer*, 6 (7): 28-35.
- Fang, Y., Wang, L., Xin, Z., Zhao, L., An, X. and Hu, Q.** (2008). Effect of foliar application of zinc, selenium, and iron fertilizers on nutrients concentration and yield of rice grain in China. *Journal of Agriculture Food and Chemistry*, 6(56): 2079-84. doi:10.1021/jf800150z
- Frossard, E., Bucher, M., Machler, F., Mozafar, A. and Hurrell, F.** (2000). Potential for increasing the content and bioavailability of Fe, Zn and Ca in plants for human nutrition. *Journal of the Science of Food and Agriculture*, 80: 861-879.
- Gannon, B., Kaliwile, C., Arscott, S. A., Schmaelzle, S., Chileshe, J., Kalungwana, N., Mosonda, M., Pixley, K., Masi, C. and Tanumihardjo S. A.** (2014). Biofortified orange maize is as efficacious as a vitamin A supplement in Zambian children even in the presence of high liver reserves of vitamin A: a community based randomized placebo-controlled trial. *American Journal of Clinical Nutrition*, DOI:10.3945/ajcn.114.087379.
- Graham, R.D., R.M. Welch, and H.E. Bouis** (2001). Addressing micronutrient malnutrition through enhancing the nutritional quality of staple foods: Principle, perspectives and knowledge gaps. *Advances in Agronomy*, 70:77-142.
- He, W., Shohag, M.J., Wei, Y., Feng, Y. and Yang, X.** (2013). Iron concentration, bioavailability, and nutritional quality of polished rice affected by different forms of foliar iron fertilizer. *Food Chemistry*, 141(4):4122-6. Doi: 10.1016/j.foodchem.2013.07.005
- Lal, R.** (2009). Soil degradation as a reason for inadequate human nutrition. *Food Security*, 1: 45-57.
- Meenakshi, J., Johnson, N., Manyong, V., Degroote, H., Javelosa, J., Yanggen, D., Naher, F., Gonzalez, C., Garcia, J. and Meng, E.** (2010). How cost-effective is biofortification in combating micronutrient malnutrition? An ex ante assessment, *World Development*, 38 (1): 64-75.
- Pfeiffer, W.H. and McClafferty, B.** (2007). *HarvestPlus: Breeding crops for better nutrition*. *Crop Science*, 47: S88-105.
- Raut, N., Sitaula, B. K. and Bajracharya, R. M.** (2010). Agricultural intensification: linking with livelihood improvement and environmental degradation in mid-hills of Nepal. *Journal of Agriculture and Environmental Sciences*, 11: 83-94.
- Shetty, P.** (2009). Incorporating nutritional considerations when addressing food insecurity. *Food Security*, 1: 431-40.
- Stein, A., Sachdev, H. and Qaim, M.** (2006). Potential impact and cost-effectiveness of Golden Rice. *Nature Biotechnology*, 24:1200-1201.
- Tanumihardjo, S. A., Anderson, C., Kaufer-Horwitz, M., Bode, L., Emenaker, N. J., Haqq, A. M., Satia, J. A., Silver H. and Stadler D. D.** (2007). Poverty, obesity and malnutrition: an international perspective recognizing the paradox. *Journal of the American Dietetic Association*, 107: 1966-1972.
- White, P.J. and Broad LeY, M.R.** (2011). Physiological limits to zinc biofortification of edible crops. *Frontier Plant Science*, 2: 80.
- Wiedenhoeft, A.C.** (2006). *Plant nutrition*. Chelsea House Publishers, USA.
- Yuan, L., Wu, L., Yang, C. and Quin, L.V.** (2013). Effects of iron and zinc foliar applications on rice plants and their grain accumulation and grain nutritional quality. *Journal of the Science of Food and Agriculture*, 93(2):254-61. doi:10.1002/jsfa.5749
- Zhu, C., Naqvi, S., Gomez-Galera, S., Pelacho, AM., Capell, T. and Christou, P.** (2007). Transgenic strategies for the nutritional enhancement of plants. *Trends in Plant Science*, 12:548-55

Zhou, C.Q., Zhang, Y.Q., Rashid, A., Ram, H., Savasli, E. and Arisoy, R.Z. (2012). Biofortification of wheat with zinc through zinc fertilization in seven countries. *Plant and Soil*, 361: 119–130.