

EFFECT OF DIFFERENT SOURCES OF NUTRIENTS ON PRODUCTIVITY AND QUALITY OF ONION (*ALLIUM CEPA L.*)

Bhavana Dhaker^{*1}, B.G. Chhipa² and R.S. Rathore³

Department of Horticulture, College of Agriculture, Career Point University, Kota, Rajasthan

² (Horticulture), Directorate of Research, MPUAT, Udaipur

³ (Horticulture), Department of Horticulture, RCA, Udaipur

Email: bhavanadhaker11@gmail.com

Received-05.07.2018, Revised-23.07.2018

Abstract: A field experiment was conducted during *Rabi* season 2016-17 to find out the effect of FYM and Vermi Compost with or without PSB and Azotobactor and rates of organic manures (50% and 100% RND) on growth attributes and yield parameters of onion (Agri found dark red) on a clay loam soil. The treatments comprised of organic, inorganic fertilizer and biofertilizers with ten treatments 100% RDF through inorganic, 100% RDF through FYM (N Basis), 100% RDF through vermicompost, 50% RDF through Inorganic Fertilizers + 50 % through FYM, 50% RDF through Inorganic Fertilizers + 50 % through vermicompost, 50% RDF through Inorganic Fertilizers + 50 % through FYM + PSB, 50% RDF through Inorganic Fertilizers + 50 % through vermicompost + PSB, 50% RDF through Inorganic Fertilizers + 25 % vermicompost + PSB, 100% RDF through FYM (N Basis) + PSB + Azotobactor and 100% RDF through vermicompost + PSB + Azotobactor. Results revealed that the application of organic manure significantly influenced the plant height (cm), number of leaves, fresh weight of leaves (g plant⁻¹) and dry weight of leaves (g plant⁻¹), diameter of bulb (cm), bulb weight (g), bulb yield (q ha⁻¹), total soluble solid (°B) and allyl propyl content (ppm) significantly increased with 100% RDF through Vermicompost + PSB + Azotobactor were significantly influenced with 100% RDF through Vermicompost + PSB + Azotobactor at 30 and 60 days of transplanting. Application of 100% RDF applied through vermicompost + PSB + Azotobactor (T₁₀) recorded maximum gross returns, net return and cost benefit ratio of onion crop.

Keywords: FYM, Growth, Onion, Vermicompost, Yield

INTRODUCTION

Onion, *Allium cepa L* is one of the important commercial bulbous vegetable crops grown throughout the world. It is a bulbous biennial herb of the most important vegetable cum condiments, spice crops demanded worldwide. India is the second largest producer of onion in the world, next to China, with 70% of the total production comes as winter crop and remaining 30% as *kharif* onion as off season crop, accounting for 11.40 per cent of the area and 10.40 per cent of the world production and 16 per cent of productivity. In India, onion is being grown in an area of 3.64 million hectares with production of 68.45 million tonnes and the average productivity is 18.82 tonnes per hectare. Maharashtra is the leading onion growing state of India (Anonymous, 2013). Organic manures not only provide plant nutrient but also improve the soil structure by effecting soil aggregates. They also decrease EC and increase water holding capacity and phosphate availability of soils, besides improving the fertilizer use efficiency and microbial activity. Bio fertilizers play a key role in increasing the availability of nutrient. Inoculation of these biofertilizers in very small quantity supplemented with sufficient amount of organic matter converts the insoluble and unavailable from of nutrient in soluble and available from of nutrients. The organic manures contain nutrients in small quantities as compared to the chemical fertilizers, also it contain growth

promoting substances like enzymes and hormones, besides improvement of soil fertility and productivity (Bhuma, 2001). Organic materials such as poultry manure, green manures and farmyard manure (FYM) can substitute for inorganic fertilizers to maintain productivity and environmental quality (Choudhary *et al.*, 2002).

MATERIALS AND METHODS

The experiment was conducted at the farmers field, village-Suwana, near Krishi Vigyan Kendra, Bhilwara, Maharana Pratap University of Agriculture & Technology, Udaipur, Rajasthan, India *Rabi* season, 2016-17 to effect of coinjoint use of organic and inorganic sources of nutrients on growth, yield and quality of onion (*Allium cepa L.*) var. Agrifound Dark Red. The soil of the experimental field was clay loam in texture and having pH 7.85, EC 0.11 dSm⁻¹, organic carbon content 0.42%, low in available N (178 kg/ha), medium P (18.7 kg/ha) and high K (348 kg/ha). The experiment was laid out in randomized block design with three replications. The treatments comprised of organic, inorganic fertilizer and biofertilizers with ten treatments T₁ -100% RDF through inorganic, T₂ - 100% RDF through FYM (N Basis), T₃ -100% RDF through vermicompost, T₄ -50% RDF through Inorganic Fertilizers + 50 % through FYM, T₅ -50% RDF through Inorganic Fertilizers + 50 % through vermicompost, T₆ -50% RDF through Inorganic

*Corresponding Author

Fertilizers + 50 % through FYM + PSB, T₇ -50% RDF through Inorganic Fertilizers + 50 % through vermicompost + PSB, T₈ -50% RDF through Inorganic Fertilizers + 25 % vermicompost + PSB, T₉ -100% RDF through FYM (N Basis) + PSB + *Azotobacter* and T₁₀ -100% RDF through vermicompost + PSB + *Azotobacter*.

The treatments of manure, chemical fertilizers and bio-fertilizers were applied as per treatment in respective plot. Vermicompost and FYM were applied prior to 15 days of transplanting of Onion. PSB and *Azotobacter* bio-fertilizer was applied at the time of transplanting. Inorganic fertilizer *i.e.* SSP and Murate of Potash were applied as a basal dose during transplanting whereas as 1/2 dose of urea was applied at the time of transplanting and remaining 1/4 – 1/4 dose applied at 30 and 60 days of transplanting respectively. The seeds of onion variety “Agrifound Dark Red” were treated with bavistin + carbendazim (SAAF) @ 3 gm kg⁻¹ before sowing in nursery. The seed of onion variety Agrifound Dark Red was raised in the nursery of 3 m long and 1.2 m wide and 10 cm above the ground level was prepared and manured as per the recommendation treatments. Treated seed were sown on 22 December 2016 in line and all the intercultural operations were done as and when required. 57 days old seedlings of uniform size were transplanted on 17 December 2017 in the prepared field. The spacing 15 cm row to row and 10 cm plant to plant was maintained. The seedlings were transplanted in cool evening according to the layout plan. A light irrigation was applied just after the transplanting and subsequent irrigation was given at an interval of 10-12 days depending upon the soil condition. Harvesting of onion was done on 31st May 2017.

The data on plant height (cm), number of leaves plant⁻¹, fresh weight of leaves, dry weight of leaves parameters were recorded at 30 and 60 DAT, whereas diameter of bulb (cm), average bulb weight (g) and bulb yield (q ha⁻¹) were recorded and thereafter, tabulated and analyzed statistically by method of analysis of variance. The diameter of bulb (cm), average bulb weight (g), bulb yield (q ha⁻¹), total soluble solid (°B) at harvest, Allyl Propyl content (ppm) and economics parameters were recorded and thereafter, tabulated and analyzed statistically by method of analysis of variance. The data were analyzed statistically and result was interpreted by using methods suggested by Panse and Sukhatme (1967).

RESULTS AND DISCUSSION

Growth parameters

The vegetative parameters like plant height (cm), number of leaves plant⁻¹, fresh weight of leaves, dry weight of leaves parameters were recorded at 30 and 60 DAT, were greatly influenced by both organic and inorganic nutrient sources. The plant height increased

significantly with the different treatments of organic manures, inorganic fertilizers and bio-fertilizer up to harvesting (Table1). This may be due to application of nutrient management through vermicompost along with bio-fertilizers, increased the photosynthetic activity, chlorophyll formation, nitrogen metabolism and auxin contents in the plants which ultimately improving the plant height. These findings were in agreement with the findings of Prabhakar *et al.* (2012). The significantly maximum leaves per plant was recorded in the treatments T₁₀ (100% RDF through Vermicompost + PSB + *Azotobacter*) at 30 and 60 days of transplanting followed by T₉ (100 % RDF through FYM + PSB + *Azotobacter*). However, the minimum leaves per plant was recorded under treatment T₂ (100 % RDF through FYM) at 30 and 60 DAT. Probable reasons for enhanced more number of leaves may be due to promotive effects of integrated nutrient management on vegetative growth which ultimately lead to more photosynthetic activities. These findings was in agreement with the findings of Jawadagi *et al.* (2012). The fresh weight of leaves was significantly influenced by various treatments of organic manures, inorganic fertilizers and bio-fertilizer. Similar results reported by Mahanthesh *et al.* (2005). Similarly, significantly maximum dry weight of leaves was exhibited in the treatment T₁₀ (100% RDF through Vermicompost + PSB + *Azotobacter*), followed by T₇ (50 % RDF through Inorganic Fertilizers + 50 % through Vermicompost + PSB). However, the minimum dry weight of leaves was observed in T₂ (100 % RDF through FYM). This may be due to application of major and micro nutrients by organic manure which increased the photosynthetic activity, chlorophyll formation, nitrogen metabolism and auxin contents in the plants which ultimately improving the dry weight of leaves.

Post-harvest parameter of crop

Diameter of bulb increased significantly with different treatments of organic manures, inorganic fertilizers and biofertilizer (Table2). This may be due to application of organic manures which provide major and micro nutrients resulted in increased the photosynthetic activity, chlorophyll formation, nitrogen metabolism and auxin contents in the plants which ultimately improving the diameter of bulb.

Significantly maximum bulb weight of bulb was exhibited in the treatment T₁₀ (100% RDF through Vermicompost + PSB + *Azotobacter*) followed by T₉ (100 % RDF through FYM + PSB + *Azotobacter*), Bulb yield per hectare differed significantly due to application of 100% RDF through Vermicompost + PSB + *Azotobacter* significantly increased the bulb yield of onion. The higher yield might be due to increase in plant height, number of leaves, and other yield attributes viz., fresh weight of whole plant, fresh and dry weight of bulb. Similar results have been reported by Shinde *et al.* (2013) and Gurjar *et al.* (2017).

The maximum TSS (12.04%) was recorded with T₁₀ followed by 11.35% TSS in T₉ (Table 2). The superior quality of onion under vermicompost treatments might be due to beneficial effect of organism which are brought about mucon deposited epidermal cell and coelomic cell of earthworm containing plant growth factor and B group vitamin. The effect of organic manure on quality parameters was also reported by Singh *et al.* (2015). The allyl propyl content (ppm) of onion influenced by different organic and inorganic treatments. Significantly maximum allyl propyl content was observed under the treatment T₁₀ (100% RDF applied through vermicompost + PSB + *Azotobacter*) and found significantly superior over rest of the treatment followed by T₉, where 100% RDF applied through FYM + PSB + *Azotobacter*. Whereas the minimum allyl propyl content at harvest was recorded in T₁ (100% RDF through inorganic fertilizer).

Economics

The cost of cultivation, gross returns, net returns and benefit cost ratio as influenced by different treatments are presented in Table 3. Application of 100% RDF applied through vermicompost + PSB + *Azotobacter* (T₁₀) recorded maximum gross returns of Rs. 153608 ha⁻¹ followed by treatment having 100% RDF applied through FYM + PSB +

Azotobacter (T₉), whereas minimum gross return (Rs 96390 ha⁻¹) was recorded in treatment T₁ (100% RDF through inorganic fertilizers). Data also revealed that the highest net return of Rs 88459 ha⁻¹ was obtained in treatment T₁₀ (100% RDF applied through vermicompost + PSB + *Azotobacter*) along with cost benefit ratio 2.36. While, lowest net return (Rs 33646 ha⁻¹) along with lowest cost benefit ratio 1.50 was observed in treatment T₂ (100 % RDF through FYM). Similar results have been reported by Gurjar *et al.* (2017).

CONCLUSION

The results of present investigation revealed plant height (cm), number of leaves, fresh weight of leaves (g plant⁻¹) and dry weight of leaves (g plant⁻¹), diameter of bulb (cm), bulb weight (g), bulb yield (q ha⁻¹), total soluble solid (°B) and allyl propyl content (ppm) were significantly influenced with 100% RDF through Vermicompost + PSB + *Azotobacter* at 30 and 60 days of transplanting. Application of 100% RDF applied through vermicompost + PSB + *Azotobacter* (T₁₀) recorded maximum gross returns, net return and cost benefit ratio.

Table 1. Effect of different organic and inorganic sources and their combinations on growth parameters of onion.

Treatment	Plant height		Number of leaves		Fresh weight of leaves (g/plant)		Dry weight of leaves (g/plant)	
	30 DAT	60 DAT	30 DAT	60 DAT	30 DAT	60 DAT	30 DAT	60 DAT
T ₁ - 100 % RDF through Inorganic Fertilizers	21.33	38.67	5.00	8.20	22.67	40.80	3.27	5.37
T ₂ - 100 % RDF through FYM (N basis)	17.40	36.67	4.80	7.84	21.67	39.00	3.20	5.00
T ₃ - 100 % RDF through Vermicompost (N basis)	19.77	37.77	5.09	8.29	26.00	46.80	3.83	5.97
T ₄ - 50 % RDF through Inorganic Fertilizers + 50 % through FYM	20.07	38.07	5.60	8.80	25.67	47.80	3.75	6.00
T ₅ - 50 % RDF through Inorganic Fertilizers + 50 % through Vermicompost	19.27	37.27	5.53	8.73	29.67	53.40	4.40	6.67
T ₆ - 50 % RDF through Inorganic Fertilizers + 50 % through FYM + PSB	22.83	43.00	5.97	9.20	28.00	47.60	4.10	5.97
T ₇ - 50 % RDF through Inorganic Fertilizers + 50 % through Vermicompost + PSB	25.47	45.13	5.87	9.13	29.67	53.40	4.45	6.62
T ₈ - 50 % RDF through Inorganic Fertilizers + 25 % Vermicompost + 25 % FYM + PSB	24.07	42.07	5.27	10.32	29.17	52.50	4.20	6.69
T ₉ - 100 % RDF through FYM + PSB + <i>Azotobacter</i>	25.00	46.33	6.20	11.34	29.50	52.30	4.32	6.83
T ₁₀ - 100 % RDF through Vermicompost + PSB + <i>Azotobacter</i>	28.67	52.00	6.93	12.63	34.33	60.53	5.05	7.69
SEm [±]	0.93	1.17	0.13	0.23	1.57	2.42	0.24	0.39
C.D. (5%)	2.77	3.47	0.38	0.67	4.66	7.20	0.72	1.15

Table 2. Effect of different organic and inorganic sources and their combinations on yield and quality parameters of onion.

Treatment	Diameter of bulb (cm)	Average weight of bulb (g)	Bulb yield (Q/ha)	Total soluble solids (°B)	Allyl propyl content (ppm)
T ₁ - 100 % RDF through Inorganic	5.78	81.00	137.70	9.27	17.00

Fertilizers					
T ₂ - 100 % RDF through FYM (N basis)	5.98	85.17	144.78	8.67	17.17
T ₃ - 100 % RDF through Vermicompost (N basis)	6.37	93.69	159.28	9.71	18.13
T ₄ - 50 % RDF through Inorganic Fertilizers + 50 % through FYM	6.64	94.53	160.71	10.05	18.69
T ₅ - 50 % RDF through Inorganic Fertilizers + 50 % through Vermicompost	7.03	104.00	176.80	10.37	19.23
T ₆ - 50 % RDF through Inorganic Fertilizers + 50 % through FYM + PSB	7.33	109.00	185.30	10.76	19.50
T ₇ - 50 % RDF through Inorganic Fertilizers + 50 % through Vermicompost + PSB	7.36	113.00	192.10	10.95	20.23
T ₈ - 50 % RDF through Inorganic Fertilizers + 25 % Vermicompost + 25 % FYM + PSB	7.20	117.17	199.18	11.11	20.80
T ₉ - 100 % RDF through FYM + PSB + <i>Azotobacter</i>	7.75	120.00	204.00	11.35	21.80
T ₁₀ - 100 % RDF through Vermicompost + PSB + <i>Azotobacter</i>	8.77	129.08	219.44	12.04	22.73
SEm \pm	0.32	1.27	2.16	0.09	0.31
C.D. (5%)	0.96	3.77	6.40	0.26	0.92

Table 3. Effect of different organic and inorganic sources and their combinations on economics of onion.

Treatment	Bulb yield (q ha ⁻¹)	Cost of cultivation (Rs. ha ⁻¹)	Gross return (Rs. ha ⁻¹)	Net return (Rs. ha ⁻¹)	BC ratio
T ₁ - 100 % RDF through Inorganic Fertilizers	137.70	57674	96390	38716	1.67
T ₂ - 100 % RDF through FYM (N basis)	144.78	67700	101346	33646	1.50
T ₃ - 100 % RDF through Vermicompost (N basis)	159.28	62699	111496	48797	1.78
T ₄ - 50 % RDF through Inorganic Fertilizers + 50 % through FYM	160.71	62687	112497	49810	1.79
T ₅ - 50 % RDF through Inorganic Fertilizers + 50 % through Vermicompost	176.80	65200	123760	58560	1.90
T ₆ - 50 % RDF through Inorganic Fertilizers + 50 % through FYM + PSB	185.30	62887	129710	66823	2.06
T ₇ - 50 % RDF through Inorganic Fertilizers + 50 % through Vermicompost + PSB	192.10	65400	134470	69070	2.06
T ₈ - 50 % RDF through Inorganic Fertilizers + 25 % Vermicompost + 25 % FYM + PSB	199.18	61637	139426	77789	2.26
T ₉ - 100 % RDF through FYM + PSB + <i>Azotobacter</i>	204.00	68150	142800	74650	2.10
T ₁₀ - 100 % RDF through Vermicompost + PSB + <i>Azotobacter</i>	219.44	65149	153608	88459	2.36

REFERENCES

Anonymous (2013). *Indian Horticulture Database-2013*, National Horticulture Board,

Ministry of Agriculture, Govt. of India, Guragaon, pp. 267.

Bhuma, M. (2001). Studies on the impact of

humic acid on sustenance of soil fertility and productivity of greengram (VBNNG-2). *M. Sc. (Agri.) Thesis*, Tamil Nadu Agricultural University, Coimbatore (India).

Choudhary, B. R., Fageria, M. S. and Dhaka, R. S. (2002). Role of growth hormones in chillies – A review. *Agric. Rev.*, 23(2): 145-148.

Gurjar, Jitendra Singh, Singh, S. S., Nagaich, K. N., Gurjar, P. K. S. and Singh, Lal. Effect of planting methods, organic, nutrient sources and bio-fertilizers on bulb yield and quality of Kharif onion (*ALLIUM CEPA L.*) *Plant Archives* Vol. 17 No. 1, 2017 pp. 439-444 ISSN 0972-5210

Jawadagi, R.S., Basavaraj, N., Patil, B.N., Naik, B.H. and Channappagoudar, B.B. (2012). Effect of different sources of nutrients on growth, yield and quality of onion (*Allium cepa L.*). *Karnataka Journal of Agricultural Sciences*. 25 (2): 232-235.

Mahanthesh, B., Venkatesha, J., Thippesha, D., Poornima, G. and Umesha, K. (2005). Effect of bio-fertilizers with levels of NPK on growth and yield of onion (*Allium cepa L.*) cv. Bellary Red grown under irrigated condition in central dry zone

of Karnataka. *Karnataka Journal of Horticulture*. 1 (3): 70-75.

Panse, V. G. and Sukhatme, P. V. (1967). Statistical Methods for Agricultural Workers, Indian Council of Agricultural Research Publication, New Delhi, pp.152-174.

Prabhakar, M., Hebbar, S.S. and Nair, A.K. (2012). Effect of organic farming practices on growth, yield and quality of rose onion (*Allium cepa*). *Indian Journal of Agricultural Sciences*. 82 (6): 500-503.

Singh, Abhishek, Ram, R. B. and Meena, M. L. (2015). Efficacy of different sources of nutrients and biofertilizers on growth yield quality of onion. *International Research Journal of Natural and Applied Sciences*, 2(10) : 64-70.

Shinde, K.G., Kadam, J.M., Bhalekar, M.N. and Pawar, P.K. (2013). Effect of organic, inorganic and biofertilizers on uptake of nutrients by onion (*Allium cepa L.*) grown under western Maharashtra conditions. *Journal of Agriculture Research and Technology*. 38 (2): 192-195.

